

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
17 January 2002 (17.01.2002)

PCT

(10) International Publication Number
WO 02/04409 A2

(51) International Patent Classification⁷: C07C 277/00

(21) International Application Number: PCT/US01/21504

(22) International Filing Date: 6 July 2001 (06.07.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/216,388 6 July 2000 (06.07.2000) US

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application:
US 60/216,388 (CIP)
Filed on 6 July 2000 (06.07.2000)

Pratt Street, Baltimore, MD 21224 (US). YU, Qian-Sheng [CN/US]; 116 Ridgefield, Lutherville, MD 21093 (US). HOLLOWAY, Harold, Wayne [US/US]; 121 Mariners Point Dr., Middle River, MD 21220 (US).

(74) Agents: MILLER, Mary, L. et al.; Needle & Rosenberg, P.C., Suite 1200, The Candler Building, 127 Peachtree Street, N.E., Atlanta, GA 30303-1811 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(71) Applicant and
(72) Inventor (*for all designated States except US*): THE GOVERNMENT OF THE UNITED STATES OF AMERICA, as represented by THE SECRETARY, DEPARTMENT OF HEALTH AND HUMAN SERVICES [US/US]; c/o National Institutes of Health, Office of, Technology Transfer, Suite 325, 6011 Executive Boulevard, Rockville, MD 20852-3804 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): GREIG, Nigel, H. [US/US]; 11 Anne Brent Garth, Phoenix, MD 21131 (US). MATTSON, Mark [US/US]; 858 E. Wheel Road, Bel Air, MD 21015 (US). ZHU, Xiaoxiang [US/US]; 6114 East

(54) Title: TETRAHYDROBENZOTHAZOLE ANALOGUES AS NEUROPROTECTIVE AGENTS

(57) Abstract: This invention relates generally to tetrahydrobenzothiazole analogues and tetrahydrobenzooxazole analogues, to pharmaceutical compositions which include these compounds and a pharmaceutically acceptable carrier, and to methods of treatment using these compounds. The invention also encompasses pharmaceutically acceptable esters, amides, and salts of such compounds. The invention further provides a method of reducing or delaying apoptosis in a population of cells, comprising contacting the population of cells with a tetrahydrobenzothiazole analogue or a tetrahydrobenzooxazole analogue, thereby reducing or delaying apoptosis in the population of cells.

WO 02/04409 A2

5

TETRAHYDROBENZOTHAZOLE ANALOGUES AS NEUROPROTECTIVE AGENTS

CROSS REFERENCE TO RELATED APPLICATION

10

This application claims priority to U. S. Provisional Application Serial No. 60/216,388, filed July 6, 2000, which is herein incorporated by reference in its entirety.

15

FIELD OF THE INVENTION

This invention relates generally to tetrahydrobenzothiazole analogues, to pharmaceutical compositions which include these compounds and a pharmaceutically acceptable carrier, and to methods of treatment using these compounds.

20

BACKGROUND OF THE INVENTION

Deposition of neurotoxic forms of amyloid β -peptide ($A\beta$) in the brain likely contributes to neuronal degeneration and dementia in Alzheimer's Disease (AD) patients. The tumor suppressor protein p53 is a key modulator of stress responses, and activation of p53 precedes apoptosis in many cell types. Moreover, up-regulation of p53 has been described as a common feature of several neurodegenerative disorders including AD, Parkinson's Disease, stroke, trauma, brain or spinal cord injury, and excitotoxic insults. Different triggers, like oxidative damage to DNA, overactivation of glutamate receptors, and disruption of cellular homeostasis, can initiate a cascade of intramolecular events that proceed via p53 activation of a death program called apoptosis. Consequently, the ability to inhibit p53 may be able to protect neurons against apoptotic insults.

30

35


In Komarov, P., et al., 285 Science 1733 (1999), pifithrin- α , a p53 inhibitor was studied for its efficiency in reducing the side effects of cancer therapy. This reference did not study novel analogues of tetrahydrobenzothiazoles. Tetrahydrobenzothiazole analogues have been synthesized in the prior art as antihelminthic compounds (anti-

5 parasitic). One method, which is incorporated by reference in its entirety, is described in Singh, A., et al., 14B Indian J. Chem. 997 (1976), which references Saldabos, I., et al., I Khim. Farm. Zh. 27 (1967), 68 Chem Abstr. 2856 (1968). In this method, tetrahydrobenzothiazole starting materials are treated with α -haloketones in a solvent.

10

SUMMARY OF THE INVENTION

In accordance with the purpose(s) of this invention, as embodied and broadly described herein, this invention, in one aspect, relates to tetrahydrobenzothiazole analogues comprising one or more analogues of Formula (I):



15


or a pharmaceutically acceptable salt or ester thereof, wherein:

X is O or S;

Y is NH, O, NR₂ or S;

20

Z is N or CH;



= is a mono-substituted, di-substituted, tri-substituted, or quad-substituted 6-member ring that is saturated or unsaturated, wherein the R₁ substituents are independently chosen from H, CN, NO₂, S, N, O, OH, COOH, halogen, and R₂, but not methyl ; and

R₂ is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member

5 selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;


10 (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of one carbon atom or from 3 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;

15 (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₂-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy, wherein the halogen is not in the para position;

(d) bisaryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and

20 (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

25 In a second aspect, the invention relates to tetrahydrobenzothiazole analogues comprising one or more analogues of formula (II):




5 or a pharmaceutically acceptable salt or ester thereof, wherein:

X is O or S;

Y is NH, O, NR₂ or S;

Z is N or CH;




= is a mono-substituted, di-substituted, tri-substituted, or quad-substituted 6-member ring that is saturated or unsaturated, wherein the R₁ substituents are independently chosen from H, CN, NO₂, S, N, O, OH, COOH, halogen, and R₂, but not methyl ; and

R₂ is selected from the group consisting of:

- 10 (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- 15 (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of one to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;
- 20 (c) aryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
- 25 (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
- 30 (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

In a third aspect, the invention relates to tetrahydrobenzothiazole analogues comprising one or more analogues of formula (III):




10 or a pharmaceutically acceptable salt or ester thereof, wherein:

R is selected from the group consisting of:

- 15 (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- 20 (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of one carbon atom or from 3 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;
- 25 (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₂-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy, wherein the halogen is not in the para position;
- (d) bisaryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl;

- 5 (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

10 In a fourth aspect, the invention relates to tetrahydrobenzothiazole analogues comprising one or more analogues of formula (IV):



or a pharmaceutically acceptable salt or ester thereof, wherein:

- 15 R is selected from the group consisting of:
- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
 - 20 (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;
 - 25 (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
 - 30 (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH,


5

halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and

10

- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

In a fifth aspect, the invention relates to tetrahydrobenzothiazole analogues comprising one or more analogues of formula (V):



15

or a pharmaceutically acceptable salt or ester thereof, wherein:

R is selected from the group consisting of:

20


- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;
- (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;

25

30

- 5 (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
- 10 (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

In a sixth aspect, the invention relates to tetrahydrobenzothiazole analogues
15 comprising one or more analogues of formula (VI):




or a pharmaceutically acceptable salt or ester thereof, wherein:

R is selected from the group consisting of:

- 20 (a) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
- (b) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
- 25 (c) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.
- 30

5

In a seventh aspect, the invention relates to tetrahydrobenzothiazole analogues comprising one or more analogues of formula (VII):



10

or a pharmaceutically acceptable salt or ester thereof, wherein:

R is selected from the group consisting of:

15


- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;
- (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
- (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and

20

25

- 5 (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

10 In an eighth aspect, the invention relates to tetrahydrobenzothiazole analogues comprising one or more analogues of formula (VIII):




15 or a pharmaceutically acceptable salt or ester thereof, wherein:

R is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- 20 (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;
- 25 (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;

- 5 (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
- 10 (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

In a ninth aspect, the invention relates to tetrahydrobenzothiazole analogues
15 comprising one or more analogues of formula (IX):




or a pharmaceutically acceptable salt or ester thereof, wherein:

R is selected from the group consisting of:

- 20 (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- 25 (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;

- 5 (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
- (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
- 10 (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.
- 15

In a tenth aspect, the invention relates to tetrahydrobenzothiazole analogues comprising one or more analogues of formula (X):



20

or a pharmaceutically acceptable salt or ester thereof, wherein:

R is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- 25 (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting
- 30

- 5 of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;
- (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
- 10 (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.
- 15

The invention also encompasses pharmaceutically acceptable esters, amides, and salts of such compounds, as will be explained in detail, *infra*.

20

Such compounds of the formula (I), (II), (III), (IV), (V), (VI), (VII), (VIII), (IX), (X), (XI) through (XV) and their pharmaceutically acceptable esters, amides, and salts are referred to herein as the inventive compounds.

25

In another aspect, the invention relates to pharmaceutical compositions containing the aforementioned inventive compounds in combination with a pharmaceutically acceptable carrier.

30 The invention further provides a method of reducing or delaying apoptosis in a population of cells, comprising contacting the population of cells with a tetrahydrobenzothiazole analogue, thereby reducing or delaying apoptosis in the population of cells.

In yet another aspect, the invention provides a method of treating a subject with a degenerative condition or of preventing a degenerative condition in a subject,

35

- 5 comprising administering to the subject a therapeutically effective amount of a tetrahydrobenzothiazole analogue.

Furthermore, the invention provides a method of treating a subject after an ischemic event to reduce ischemia-induced apoptosis, comprising administering to the
10 subject a therapeutically effective amount of a tetrahydrobenzothiazole analogue, thereby reducing apoptosis.

Additional advantages of the invention will be set forth in part in the description which follows, and in part will be obvious from the description, or may be learned by
15 practice of the invention. The advantages of the invention will be realized and attained by means of the elements and combinations particularly pointed out in the appended claims. It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory only and are not restrictive of the invention, as claimed.

20

The accompanying drawings, which are incorporated in and constitute a part of this specification, illustrate several embodiments of the invention and together with the description, serve to explain the principles of the invention.

25

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 provides the chemical structures and data for the compounds tested in examples 4 and 5.

30 Figure 2 shows the effect PFT α (compound 5), its precursor, or analogues thereof on PC12 cells treated with the DNA-damaging agent camptothecin. PC12 cells were pretreated for 6 h with compounds (precursor, 1, 4-11, 13-15) prepared in 0.5% DMSO at concentrations between 100-400 nM) and were then exposed for 24 hours to camptothecin (40 μ M). Figure 2A shows the amount of fluorescence emitted by the
35 live-cell indicator dye, 2', 7'-bis(2-carboxyethyl)-5-6-carboxyfluorescein AM ester (BCECF AM: 5 μ M). This fluorescent dye was taken up and retained by live cells.

5 Figure 2B shows the quantification of neuronal survival in the cultures (n=3 cultures). PFT α (compound 5) and several of its analogues protected cultured PC12 cells against death induced by camptothecin.

Figure 3 shows the results of a neuronal survival assay using cultured
10 hippocampal neurons treated with either camptothecin or etoposide in the presence and absence of PFT α (compound 5) or its precursor. PFT α protects cultured hippocampal neurons against death induced by DNA-damaging agents camptothecin and etoposide. Hippocampal cell cultures were pretreated for 1 hour with 0.5% DMSO (vehicle), 200 nM PFT α or 200 nM of PFT α precursor. Cultures were then exposed for 24 hours to
15 the DNA-damaging agents camptothecin (5 μ M) and etoposide (2.5 μ M). Neuronal survival in each culture was quantified (n=4-6 cultures). **p<0.01 compared to corresponding value for cultures treated with vehicle or precursor (ANOVA with Scheffe post-hoc tests).

Figure 4 shows the results of a neuronal survival assay using cultured
20 hippocampal neurons treated with either camptothecin or etoposide in the presence and absence of PFT α and its analogues 14, 1, 15, 5x. Compounds were added 1h before exposure of primary hippocampal neurons to the DNA-damaging compounds camptothecin (5 μ M) or etoposide (2.5 μ M). The percentage of neuronal survival 24h after the treatment is given for each group as mean \pm SD.

25 Figure 5 shows the results of a neuronal survival assay using cultured hippocampal neurons treated with camptothecin in the presence and absence of PFT α analogues 4, 6-8, 9-11, 13, 16 (100nM). Compounds were added 1h before exposure of primary hippocampal neurons to the DNA-damaging compound camptothecin (5 μ M). Compounds 8 and 9 significantly protected against toxicity (p \leq 0.05).

30 Figure 6 shows the results of histologic analysis of four different brain levels in mice following transient focal cerebral ischemia with and without pretreatment with PFT- α (compound 5). PFT- α (2mg/kg) was administered intraperitoneally 1h before transient middle cerebral artery occlusion in C57BL/6 mice. Twenty four hours after reperfusion, mice were euthanized and the infarct size was quantified after TTC-staining of 2mm brain sections. Figure 2A shows the infarct area in mm² at each level.
35

- 5 Figure 2B shows the infarct volume in mm^3 . Values are the mean and SD of 12 animals per group.

Figure 7 shows the results of behavioral tests following PFT- α and compounds 5x and 13 treatment in a MPTP-induced Parkinson's disease model. C57BL/6 mice
10 were given vehicle (control) or MPTP, and PFT-alpha (2mg/kg), compound 5x (2mg/kg), compound 13 (2mg/kg), or vehicle were administered 30 minutes before the first MPTP injection and again 30 minutes after the last MPTP injection. Mice were tested on a rotarod apparatus. Figure 7A shows the summary data for measurements of numbers of falls. Figure 7B shows the summary data for running times. Values are the
15 mean and SE (standard error) of determinations made in 12 mice/group. (* $P < 0.01$ compared to control value, ** $P < 0.01$ compared to MPTP value. ANOVA with Scheffe post-hoc test).

Figure 8 shows the results of densitometric analysis of Western blots stained
20 with a TH antibody. PFT-alpha and compound 5x attenuate MPTP-induced loss of striatal tyrosine hydroxylase. Mice were given saline (control) or MPTP, and PFT-alpha (2mg/kg), compound 5x (2mg/kg), compound 13 (2mg/kg) or vehicle were administered 30 minutes before the first MPTP injection and were repeated 30 minutes after the last MPTP injection. 7 days later, mice were euthanatized and striatal tissue
25 samples were removed. Levels of TH were determined by immunoblot analysis (Western blots). The results of densitometric analysis of blots of samples from 4-6 different mice/group are shown as mean \pm SE. (* $P < 0.01$ compared to control value, ** $P < 0.01$ compared to MPTP value. ANOVA with Scheffe post-hoc tests).

30 Figure 9 shows the results of quantification of TH-positive cells in the substantia nigra of animals treated with MPTP, with or without PFT- α and compounds 5x and 13. PFT- α and compound 5x attenuated MPTP-induced loss of substantia nigra dopaminergic neurons. Mice were given saline (control) or MPTP, and PFT-alpha (2mg/kg) (compound 5), Z-1-117 (2mg/kg) (compound 5x), Z-1-143 (2mg/kg)
35 (compound 13) or vehicle were administered 30 minutes before the first MPTP injection and were repeated 30 minutes after the last MPTP injection. Values are shown

- 5 the mean \pm SE of counts made in 4 mice/group. (*P<0.05 , **P<0.01 compared to MPTP value. ANOVA with Scheffe post-hoc tests).

DESCRIPTION OF THE PREFERRED EMBODIMENTS

10 The present invention may be understood more readily by reference to the following detailed description of preferred embodiments of the invention and the Examples included therein and to the Figures and their previous and following description.

15 Before the present compounds, compositions, articles, devices, and/or methods are disclosed and described, it is to be understood that this invention is not limited to specific synthetic methods of using or making as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting.

20 It must be noted that, as used in the specification and the appended claims, the singular forms "a," "an" and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "an aromatic compound" includes mixtures of aromatic compounds, reference to "a pharmaceutical carrier" includes mixtures of two or more such carriers, and the like.

25 Ranges may be expressed herein as from "about" one particular value, and/or to "about" another particular value. When such a range is expressed, another embodiment includes from the one particular value and/or to the other particular value. Similarly, when values are expressed as approximations, by use of the antecedent "about," it will be understood that the particular value forms another embodiment. It will be further understood that the endpoints of each of the ranges are significant both in relation to the other endpoint, and independently of the other endpoint.

35 In this specification and in the claims which follow, reference will be made to a number of terms which shall be defined to have the following meanings:

5 References in the specification and concluding claims to parts by weight, of a particular element or component in a composition or article, denotes the weight relationship between the element or component and any other elements or components in the composition or article for which a part by weight is expressed. Thus, in a compound containing 2 parts by weight of component X and 5 parts by weight
10 component Y, X and Y are present at a weight ratio of 2:5, and are present in such ratio regardless of whether additional components are contained in the compound.

 A weight percent of a component, unless specifically stated to the contrary, is based on the total weight of the formulation or composition in which the component is
15 included.

 A residue of a chemical species, as used in the specification and concluding claims, refers to the moiety that is the resulting product of the chemical species in a particular reaction scheme or subsequent formulation or chemical product, regardless of
20 whether the moiety is actually obtained from the chemical species. Thus, an ethylene glycol residue in a polyester refers to one or more $\text{-OCH}_2\text{CH}_2\text{O-}$ units in the polyester, regardless of whether ethylene glycol was used to prepare the polyester. Similarly, a sebacic acid residue in a polyester refers to one or more $\text{-CO(CH}_2)_8\text{CO-}$ moieties in the polyester, regardless of whether the residue is obtained by reacting sebacic acid or an
25 ester thereof to obtain the polyester.

 The term "halogen" and "halo" refer to bromine, chlorine, fluorine, and iodine.

 The term "alkyl" as used herein refers to a branched or unbranched saturated
30 hydrocarbon group, such as methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, isobutyl, *t*-butyl, octyl, decyl, tetradecyl, hexadecyl, eicosyl, tetracosyl and the like. The term "lower alkyl" intends an alkyl group of from one to six carbon atoms, preferably from one to four carbon atoms. The term "cycloalkane" as used herein refers to a cyclic alkane group.

35

5 The term "alkoxy" as used herein intends an alkyl group bound through a single, terminal ether linkage; that is, an "alkoxy" group may be defined as -OR where R is alkyl as defined above. A "lower alkoxy" group intends an alkoxy group containing from one to six, more preferably from one to four, carbon atoms.

10 The term "alkylene" as used herein refers to a difunctional saturated branched or unbranched hydrocarbon chain, for example, methylene (-CH₂-), ethylene (-CH₂-CH₂-), propylene (-CH₂-CH₂-CH₂-), 2-methylpropylene [-CH₂-CH(CH₃)-CH₂-], hexylene [-CH₂]₆- and the like. "Lower alkylene" refers to an alkylene group of from 1 to 6, more preferably from 1 to 4, carbon atoms. The term "cycloalkylene" as used herein
15 refers to a cyclic alkylene group.

 The term "alkene" as used herein intends a mono-unsaturated or di-unsaturated hydrocarbon group. Asymmetric structures such as (AB)C=C(CD) are intended to include both the *E* and *Z* isomers. This may be presumed in structural formulae herein
20 wherein an asymmetric alkene is present, or it may be explicitly indicated by the bond symbol —.



 The term "aryl" as used herein refers to a C₆H₆ aromatic ring. Substituents on the aryl group may be present on any position, i.e. *ortho*, *meta* or *para* positions or
25 fused to the aromatic ring.

 By "bisphenol" it is meant that two C₆H₆ aromatic rings are present in a group in an unfused state. The aromatic rings may be joined at any position, i.e. *ortho*, *meta* or *para* positions, relative to the attachment position to the structure. Substituents on
30 the bisphenol group may be present on or fused to any position of either aromatic ring.

 The term "condensed aromatic" as used herein refers to more than one fused C₆H₆ aromatic ring. The aromatic rings may be fused at any bond i.e. along the C₂-C₃ bond relative to the C₁ attachment position to the structure. Substituents on the
35 condensed aromatic group may be present on or fused to any position of any of the aromatic rings.

5

By "tetrahydrobenzothiazole" or "tetrahydrobenzothiazole analogue" it is meant to include tetrahydrobenzothiazole analogues, tetrahydrobenzooxazole analogues, and heterocyclic analogues of the general formulas (GF1) and (GF2):



10

(GF1)


(GF2)

or a pharmaceutically acceptable salt or ester thereof, wherein:

X is O or S;

Y is NH, O, NR₂ or S;

Z is N or CH;



= is a mono-substituted, di-substituted, tri-substituted, or quad-substituted 6-member ring that is saturated or unsaturated, wherein the R₁ substituents are independently chosen from H, CN, NO₂, S, N, O, OH, COOH, halogen, and R₂; and

15

R₂ is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH and halogen;
- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;


20

25

- 5 (c) aryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
- 10 (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
- 15 (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

By the terms "PFT α ," "PFT-a," and "pififthr α " is meant [2-(2-imino-4,5,6,7-tetrahydrobenzothiazol-3-yl)-1-p-tolyethanone] which is represented by the chemical structure (PFT α):

20



5

(PFT α)

or a pharmaceutically acceptable salt or ester thereof.

By "substituted" is meant that the substituent group may be present anywhere
10 within or attached to the substituted group.

By "neural cell" is meant any cell that can be located in the central or peripheral nervous system or is a precursor or derivative thereof, including, for example, neuronal cells, glial cells, neural stem cells, neuronal stem cells, neuroblasts. By "cardiac cell"
15 is meant any cell that can be located in the cardiac tissue or is a precursor or derivative thereof, including, for example, cardiomyocytes, cardiac stem cells, endothelial cells, and myoblasts. By "pancreatic islet cell" is meant any cells type present in the pancreatic islet or precursors or derivatives thereof, including, for example, alpha cells (glucagon secreting cells) and beta cells (insulin secreting cells). By "muscle cells" is
20 meant skeletal, smooth, or cardiac muscle cells or precursors or derivatives thereof, including, for example, myoblasts. Any of these cell populations can include immortalized cells and can include transfected or transformed cells. Thus, the population of neural cells, cardiac cells, pancreatic cells, or muscle cells can include one or more types of neural cells, cardiac cells, pancreatic cells, or muscle cells.

25

By "reducing or delaying" is meant either slowing or eliminating all or a portion of the apoptosis so that cell death is diminished or delayed in one or more cells in the cellular population.

30 As used throughout, by "contacting" is meant an instance of exposure of at least one cell (e.g., a neural cell, a stem cell, a cardiac cell) to an agent (e.g., a tetrahydrobenzothiazole analogue).

As used herein, "a degenerative condition" means a disease or condition marked
35 by cell death. The degenerative condition can include, for example, neurodegenerative diseases (e.g., Alzheimer's disease, Parkinson's disease, Huntington's disease,

5 amyotrophic lateral sclerosis, multiple sclerosis, brain and spinal cord injury, peripheral neuropathies, and stroke), degenerative cardiomyopathies (e.g., idiopathic dilated, ischemic, hypertrophic, obstructive, familial obstructive, familial arrhythmogenic right, ventricular, post-viral, alcoholic, endomyocardial fibrosis, amyloidosis, and muscular dystrophy), degenerative myopathies (e.g., muscular dystrophies), and other
10 degenerative processes (e.g., diabetes Type I or Type II, and ischemia).

As used throughout, by "subject" is meant an individual. Preferably, the subject is a mammal such as a primate, and, more preferably, a human. Thus, the "subject" can include domesticated animals, such as cats, dogs, etc., livestock (e.g., cattle, horses,
15 pigs, sheep, goats, etc.), and laboratory animals (e.g., mouse, rabbit, rat, guinea pig, etc.).

"Optional" or "optionally" means that the subsequently described event or circumstance may or may not occur, and that the description includes instances where
20 said event or circumstance occurs and instances where it does not. For example, the phrase "optionally substituted lower alkyl" means that the lower alkyl group may or may not be substituted and that the description includes both unsubstituted lower alkyl and lower alkyl where there is substitution.

25 In general, "a therapeutically effective dose or doses" means the amount needed to achieve the desired result or results (reducing or delaying apoptosis or treating a degenerative condition). One of ordinary skill in the art will recognize that the potency and, therefore, a "therapeutically effective dose or doses" can vary for the various tetrahydrobenzothiazole analogues used in this invention. One skilled in the art can
30 readily assess the potency of the analogues.

The term "modified" is often used herein to describe polymers and means that a particular monomeric unit that would typically make up the pure polymer has been replaced by another monomeric unit that shares a common polymerization capacity
35 with the replaced monomeric unit. Thus, for example, it is possible to substitute diol residues for glycol in poly(ethylene glycol), in which case the poly(ethylene glycol)

5 will be "modified" with the diol. If the poly(ethylene glycol) is modified with a mole percentage of the diol, then such a mole percentage is based upon the total number of moles of glycol that would be present in the pure polymer but for the modification. Thus, in a poly(ethylene glycol) that has been modified by 50 mole % with a diol, the diol and glycol residues are present in equimolar amounts.

10


By "pharmaceutically acceptable" is meant a material that is not biologically or otherwise undesirable, i.e., the material may be administered to an individual along with the selected bicyclic compound without causing any undesirable biological effects or interacting in a deleterious manner with any of the other components of the pharmaceutical composition in which it is contained.

15

The invention also encompasses pharmaceutically acceptable nontoxic ester, amide, and salt derivatives of those compounds of formulas (I), (II), (III), (IV), (V), (VI), (VII), (VIII), (IX), and (X) containing a carboxylic acid moiety.

20

In one embodiment, the tetrahydrobenzothiazole analogue comprises one or more analogues of Formula (I):



(I)

25


or a pharmaceutically acceptable salt or ester thereof, wherein:

X is O or S;

Y is NH, O, NR₂ or S;

5

Z is N or CH;



= is a mono-substituted, di-substituted, tri-substituted, or quad-substituted 6-member ring that is saturated or unsaturated, wherein the R_1 substituents are independently chosen from H, CN, NO₂, S, N, O, OH, COOH, halogen, and R_2 , but not methyl; and

R_2 is selected from the group consisting of:

10

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, preferably from 1 to 8 carbon atoms, more preferably from 1 to four carbon atoms, wherein the carbon chain is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;

15

- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of one carbon atom or from 3 to 20 carbon atoms, preferably one or 3 to 8 carbon atoms, more preferably one, three or four carbon atoms, wherein the carbon chain is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl; preferably substituted with at least one member selected from the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more preferably substituted with at least one member selected from the group consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl;

20


25

- (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₂-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy; preferably substituted with at least one member selected from the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more preferably substituted with at least one member selected from the group consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl; wherein the halogen is not in the para position;

30


- 5 (d) bisaryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; preferably substituted with at least one member selected from the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more preferably substituted with at least one member selected from the group consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl; and
- 10 (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; preferably substituted with at least one member selected from the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more preferably substituted with at least one member selected from the group consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl.

In another embodiment, the tetrahydrobenzothiazole analogue comprises one or
 20 more analogues of formula (II):



or a pharmaceutically acceptable salt or ester thereof, wherein:

- 25 X is O or S;
 Y is NH, O, NR₂ or S;
 Z is N or CH;



= is a mono-substituted, di-substituted, tri-substituted, or quad-substituted 6-member ring that is saturated or unsaturated, wherein the R₁ substituents are independently chosen from H, CN, NO₂, S, N, O, OH, COOH, halogen, and R₂, but not methyl ; and

5

R₂ is selected from the group consisting of:

10

(a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, preferably from 1 to 8 carbon atoms, more preferably from 1 to four carbon atoms, wherein the carbon chain is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;

15

(b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of one to 20 carbon atoms, preferably one to 8 carbon atoms, more preferably one to four carbon atoms, wherein the carbon chain is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl; preferably substituted with at least one member selected from the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more preferably substituted with at least one member selected from the group consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl;

20

25

(c) aryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy; preferably substituted with at least one member selected from the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more preferably substituted with at least one member selected from the group consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl;


30

35

(d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; preferably substituted with at least one member selected from the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more preferably substituted with at least one member selected from the group consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl; and

- 5 (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; preferably substituted with at least one member selected from the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more preferably substituted with at least one member selected from the group consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl.
- 10

In a further embodiment, the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (III):



or a pharmaceutically acceptable salt or ester thereof, wherein:

R is selected from the group consisting of:


- 20 (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, preferably from 1 to 8 carbon atoms, more preferably from 1 to four carbon atoms, wherein the carbon chain is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- 25 (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of one carbon atom or from 3 to 20 carbon atoms, preferably one or 3 to 8 carbon atoms, more preferably one, three or four carbon atoms, wherein the carbon chain is unsubstituted or substituted with at least one member selected from the group consisting of CN,

- 5 NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl; preferably substituted with at least one member selected from the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more preferably substituted with at least one member selected from the group consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl;
- 10 (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₂-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy; preferably substituted with at least one member selected from the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more preferably substituted with at least one member selected from the group consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl; wherein the halogen is not in the para position;
- 15 (d) bisaryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; preferably substituted with at least one member selected from the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more preferably substituted with at least one member selected from the group consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl; and
- 20 (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; preferably substituted with at least one member selected from the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more preferably substituted with at least one member selected from the group consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl.
- 25
- 30

Some preferred embodiments of formula (III) include where R is ethyl, methoxy, methoxyphenyl, fluorophenyl, chlorophenyl, nitrophenyl, or

35 tetradecyloxyphenyl.

5 Moreover, in another embodiment, the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (IV):



or a pharmaceutically acceptable salt or ester thereof, wherein:

10 R is selected from the group consisting of:


- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, preferably from 1 to 8 carbon atoms, more preferably from 1 to four carbon atoms, wherein the carbon chain is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- 15
- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of one to 20 carbon atoms, preferably one to 8 carbon atoms, more preferably one to four carbon atoms, wherein the carbon chain is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl; preferably substituted with at least one member selected from the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more preferably substituted with at least one member selected from the group consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl;
- 20
- (c) aryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy; preferably substituted with at least one member selected from
- 25
- 30

- 5 the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more preferably substituted with at least one member selected from the group consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl;
- 10 (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; preferably substituted with at least one member selected from the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more preferably substituted with at least one member selected from the group consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl; and
- 15 (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; preferably substituted with at least one member selected from the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more preferably substituted with at least one member selected from the group consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl.
- 20

Preferred embodiments of formula (IV) include where R is ethyl, methoxy, methoxyphenyl, fluorophenyl, chlorophenyl, nitrophenyl, or tetradecyloxyphenyl.

25

In still a further embodiment, the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (V):



5

or a pharmaceutically acceptable salt or ester thereof, wherein:


R is selected from the group consisting of:

- 10 (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, preferably from 1 to 8 carbon atoms, more preferably from 1 to four carbon atoms, wherein the carbon chain is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- 15 (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of one to 20 carbon atoms, preferably one to 8 carbon atoms, more preferably one to four carbon atoms, wherein the carbon chain is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl; preferably substituted with at least one member selected from the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more preferably substituted with at least one member selected from the group consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl;
- 20 (c) aryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy; preferably substituted with at least one member selected from the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more preferably substituted with at least one member selected from the group consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl;
- 25 (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; preferably substituted with at least one member selected from the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more
- 30
- 35

- 5 preferably substituted with at least one member selected from the group consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl; and
- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; preferably substituted with at least one member selected from the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more preferably substituted with at least one member selected from the group consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl.

15 Preferred embodiments of formula (V) include where R is methyl, benzyl, cyclopropylmethyl, fluorobenzyl, difluorobenzyl, chlorobenzyl, dichlorobenzyl, methylbenzyl, t-butylbenzyl, nitrobenzyl, cyanobenzyl, trifluoromethylbenzyl, phenylbenzyl, menaphthyl, phenylethyl, or butyl.

20 In another preferred embodiment, the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (VI):




or a pharmaceutically acceptable salt or ester thereof, wherein:

- 25 R is selected from the group consisting of:
- (a) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy; preferably substituted with at least one member selected from the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more preferably substituted
- 30

- 5 with at least one member selected from the group consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl;
- (b) bisaryl which is unsubstituted or substituted with at least one member
10 selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; preferably substituted with at least one member selected from the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more preferably substituted with at least one member selected from the group consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl; and
- (c) condensed aromatic which is unsubstituted or substituted with at least
15 one member selected from the group consisting CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; preferably substituted with at least one member selected from the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more preferably substituted with at least one member selected
20 from the group consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl.

Preferred embodiments of formula (VI) include where R is methyl, benzyl, cyclopropylmethyl, fluorobenzyl, difluorobenzyl, chlorobenzyl, dichlorobenzyl, methylbenzyl, t-butylbenzyl, nitrobenzyl, cyanobenzyl, trifluoromethylbenzyl,
25 phenylbenzyl, menaphthyl, phenylethyl, or butyl.

In another preferred embodiment, the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (VII):



5

or a pharmaceutically acceptable salt or ester thereof, wherein:

R is selected from the group consisting of:


- 10 (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, preferably from 1 to 8 carbon atoms, more preferably from 1 to four carbon atoms, wherein the carbon chain is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- 15 (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of one to 20 carbon atoms, preferably one to 8 carbon atoms, more preferably one to four carbon atoms, wherein the carbon chain is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl; preferably substituted with at least one member selected from the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more preferably substituted with at least one member selected from the group consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl;
- 20 (c) aryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy; preferably substituted with at least one member selected from the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more preferably substituted with at least one member selected from the group consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl;
- 25 (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; preferably substituted with at least one member selected from the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more
- 30
- 35

- 5 preferably substituted with at least one member selected from the group consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl; and
- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; preferably substituted with at least one member selected from the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more preferably substituted with at least one member selected from the group consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl.

15 Preferred embodiments of formula (VII) include where R is methyl, benzyl, cyclopropylmethyl, fluorobenzyl, difluorobenzyl, chlorobenzyl, dichlorobenzyl, methylbenzyl, t-butylbenzyl, nitrobenzyl, cyanobenzyl, trifluoromethylbenzyl, phenylbenzyl, menaphthyl, phenylethyl, or butyl.

20 A preferred embodiment of the present invention includes the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (VIII):

(VIII)




or a pharmaceutically acceptable salt or ester thereof, wherein:

- 25 R is selected from the group consisting of:
- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, preferably from 1 to 8 carbon atoms, more preferably from 1 to four carbon atoms, wherein the carbon chain is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- 30

- 5 (b) alkoxy which is straight chain, branched chain or cyclic, having a
carbon chain length of one to 20 carbon atoms, preferably one to 8
carbon atoms, more preferably one to four carbon atoms, wherein the
carbon chain is unsubstituted or substituted with at least one member
selected from the group consisting of CN, NO₂, S, N, O, OH, COOH,
10 halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl; preferably
substituted with at least one member selected from the group consisting
of OH, COOH, halogen, and C₁-C₈ alkyl; more preferably substituted
with at least one member selected from the group consisting of OH,
COOH, Cl, F, Br, and C₁-C₄ alkyl;
- 15 (c) aryl which is unsubstituted or substituted with at least one member
selected from the group consisting of CN, NO₂, S, N, O, OH, COOH,
halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀
alkoxy; preferably substituted with at least one member selected from
the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more
20 preferably substituted with at least one member selected from the group
consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl;
- (d) bisaryl which is unsubstituted or substituted with at least one member
selected from the group consisting of CN, NO₂, S, N, O, OH, COOH,
halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy,
25 and aryl; preferably substituted with at least one member selected from
the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more
preferably substituted with at least one member selected from the group
consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl; and
- (e) condensed aromatic which is unsubstituted or substituted with at least
30 one member selected from the group consisting CN, NO₂, S, N, O, OH,
COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀
alkoxy, and aryl; preferably substituted with at least one member
selected from the group consisting of OH, COOH, halogen, and C₁-C₈
alkyl; more preferably substituted with at least one member selected
35 from the group consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl.

5 Preferred embodiments of formula (VIII) include where R is methyl, benzyl, cyclopropylmethyl, fluorobenzyl, difluorobenzyl, chlorobenzyl, dichlorobenzyl, methylbenzyl, t-butylbenzyl, nitrobenzyl, cyanobenzyl, trifluoromethylbenzyl, phenylbenzyl, menaphthyl, phenylethyl, or butyl.

10 In another preferred embodiment, the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (IX):



or a pharmaceutically acceptable salt or ester thereof, wherein:

15 R is selected from the group consisting of:


- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, preferably from 1 to 8 carbon atoms, more preferably from 1 to four carbon atoms, wherein the carbon chain is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- 20
- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of one to 20 carbon atoms, preferably one to 8 carbon atoms, more preferably one to four carbon atoms, wherein the carbon chain is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl; preferably substituted with at least one member selected from the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more preferably substituted
- 25

- 5 with at least one member selected from the group consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl;
- (c) aryl which is unsubstituted or substituted with at least one member
selected from the group consisting of CN, NO₂, S, N, O, OH, COOH,
halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀
10 alkoxy; preferably substituted with at least one member selected from
the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more
preferably substituted with at least one member selected from the group
consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl;
- (d) bisaryl which is unsubstituted or substituted with at least one member
15 selected from the group consisting of CN, NO₂, S, N, O, OH, COOH,
halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy,
and aryl; preferably substituted with at least one member selected from
the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more
preferably substituted with at least one member selected from the group
20 consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl; and
- (e) condensed aromatic which is unsubstituted or substituted with at least
one member selected from the group consisting CN, NO₂, S, N, O, OH,
COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀
alkoxy, and aryl; preferably substituted with at least one member
25 selected from the group consisting of OH, COOH, halogen, and C₁-C₈
alkyl; more preferably substituted with at least one member selected
from the group consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl.

Preferred embodiments of formula (IX) include where R is methyl, benzyl,
30 cyclopropylmethyl, fluorobenzyl, difluorobenzyl, chlorobenzyl, dichlorobenzyl,
methylbenzyl, t-butylbenzyl, nitrobenzyl, cyanobenzyl, trifluoromethylbenzyl,
phenylbenzyl, menaphthyl, phenylethyl, or butyl.

5

In another preferred embodiment, the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (X):



10

or a pharmaceutically acceptable salt or ester thereof, wherein:

R is selected from the group consisting of:

15

(a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, preferably from 1 to 8 carbon atoms, more preferably from 1 to four carbon atoms, wherein the carbon chain is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;

20

(b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of one to 20 carbon atoms, preferably one to 8 carbon atoms, more preferably one to four carbon atoms, wherein the carbon chain is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl; preferably substituted with at least one member selected from the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more preferably substituted with at least one member selected from the group consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl;

25

30

(c) aryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy; preferably substituted with at least one member selected from

- 5 the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more preferably substituted with at least one member selected from the group consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl;
- 10 (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; preferably substituted with at least one member selected from the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more preferably substituted with at least one member selected from the group consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl; and
- 15 (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; preferably substituted with at least one member selected from the group consisting of OH, COOH, halogen, and C₁-C₈ alkyl; more preferably substituted with at least one member selected
- 20 from the group consisting of OH, COOH, Cl, F, Br, and C₁-C₄ alkyl.

Preferred embodiments of formula (X) include where R is methyl, benzyl, cyclopropylmethyl, fluorobenzyl, difluorobenzyl, chlorobenzyl, dichlorobenzyl, 25 methylbenzyl, t-butylbenzyl, nitrobenzyl, cyanobenzyl, trifluoromethylbenzyl, phenylbenzyl, menaphthyl, phenylethyl, or butyl.

In a preferred embodiment for each of the tetrahydrobenzothiazole analogues having the formula (I), (II), (III), (IV), (V), (VII), (VIII), (IX), or (X), subparts (b), (c), 30 (d), and (e) or formula (VI), subparts (a), (b) and (c) may preferably be C₁-C₈ alkenyl or C₁-C₈ alkynyl, more preferably C₁-C₄ alkenyl or C₁-C₄ alkynyl.

In one embodiment of the method of reducing or delaying apoptosis, the tetrahydrobenzothiazole analogue is pifithrin- α . Alternatively, the 35 tetrahydrobenzothiazole analogue is selected from the group of analogues having formula (I), (II), (III), (IV), (V), (VI), (VII), (VIII), (IX), (X) or (XI) through (XV).

5

The present method is useful in reducing or delaying apoptosis induced by a toxin (e.g., the neurotoxic form of amyloid β -peptide, camptothecin, glutamate, anti-cancer drugs, such as etoposide and derivatives thereof; vinca alkaloids; and chemical agents, such as 3-nitropropionic acid, 1-methyl-4-phenyl-1, 2, 3, 6-tetrahydropyridine (MPTP), domoic acid, and kainic acid), ischemia (e.g., caused by a stroke, transient ischemic attack, myocardial infarction), trauma (e.g., head injury), genetic defect (e.g., mutations in amyloid precursor protein; presenilins; α -synuclein; copper, zinc superoxide dismutase; Notch3 gene), genetic disease (e.g., muscular dystrophy, Huntington's disease, CADASIL (cerebral autosomal dominant arteriopathy with subcortical infarcts and leukoencephalopathy), environmental factors, (e.g., irradiation or severe seizure disorders).

Also provided is a method of treating a subject with a degenerative condition or of reducing one or more symptoms of a degenerative condition in a subject, comprising administering to the subject a therapeutically effective amount of a tetrahydrobenzothiazole analogue. The degenerative condition may result from events such as, but not limited to, surgery or organ transplant, trauma, severe seizure disorder, environmental factors, toxins, ischemia, genetic defects or diseases. With such conditions, one embodiment of the present invention provides a method of treating a subject before a surgical procedure to reduce apoptosis, comprising administering to the subject a therapeutically effective amount of a tetrahydrobenzothiazole analogue, thereby reducing apoptosis. In another embodiment, the invention provides a method of treating a subject after an excitotoxic event. Such methods are also useful in treating a subject before a trigger event such as an ischemic or excitotoxic event when such an event can be foreseen. In another embodiment, the invention provides a method of treating a subject after an ischemic event to reduce ischemia-induced apoptosis, comprising administering to the subject a therapeutically effective amount of a tetrahydrobenzothiazole analogue, thereby reducing apoptosis.

The invention also provides a method of treating a subject exposed to irradiation, comprising administering to the subject a therapeutically effective amount

5 of a tetrahydrobenzothiazole analogue, thereby reducing one or more of the undesired symptoms of irradiation. In one embodiment, the irradiation is caused by radiation therapy. The irradiating event, in another embodiment, results in radiation poisoning. Thus, the analogue is administered before, during, or after the irradiating event.

10 Pharmaceutically acceptable salts are prepared by treating the free acid with an appropriate amount of a pharmaceutically acceptable base. Representative pharmaceutically acceptable bases are ammonium hydroxide, sodium hydroxide, potassium hydroxide, lithium hydroxide, calcium hydroxide, magnesium hydroxide, ferrous hydroxide, zinc hydroxide, copper hydroxide, aluminum hydroxide, ferric
15 hydroxide, isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, ethanolamine, 2-dimethylaminoethanol, 2-diethylaminoethanol, lysine, arginine, histidine, and the like. The reaction is conducted in water, alone or in combination with an inert, water-miscible organic solvent, at a temperature of from about 0°C to about 100°C, preferably at room temperature. The molar ratio of
20 compounds of structural formula (I) to base used are chosen to provide the ratio desired for any particular salts. For preparing, for example, the ammonium salts of the free acid starting material—a particular preferred embodiment—the starting material can be treated with approximately one equivalent of pharmaceutically acceptable base to yield a neutral salt. When calcium salts are prepared, approximately one-half a molar
25 equivalent of base is used to yield a neutral salt, while for aluminum salts, approximately one-third a molar equivalent of base will be used.

Ester derivatives are typically prepared as precursors to the acid form of the compounds, as illustrated in the examples below, and accordingly may serve as
30 prodrugs. Generally, these derivatives will be lower alkyl esters such as methyl, ethyl, and the like. Amide derivatives $-(CO)NH_2$, $-(CO)NHR$ and $-(CO)NR_2$, where R is lower alkyl, may be prepared by reaction of the carboxylic acid-containing compound with ammonia or a substituted amine.

5

Synthetic Methods:

The compounds of the invention may be readily synthesized using techniques generally known to synthetic organic chemists. Suitable experimental methods for making and derivatizing aromatic compounds are described, for example, in the references cited in the Background section herein above, the disclosures of which are hereby incorporated by reference for their general teachings and for their synthesis teachings. Methods for making specific and preferred compounds of the present invention are described in detail in the Examples.

15 Utility and Administration:

The compounds of the invention defined by structural formulas (I), (II), (III), (IV), (V), (VI), (VII), (VIII), (IX), (X), (XI) through (XV), including the pharmacologically acceptable esters, amides or salts thereof, are useful in reducing or delaying apoptosis in a population of cells, comprising contacting the population of cells with a tetrahydrobenzothiazole analogue, thereby reducing or delaying apoptosis in the population of cells. In one embodiment, the cells are neural cells, and in another embodiment, the cells are cardiac cells. In another embodiment, the cells are pancreatic islet cells, and in yet another embodiment, the cells are muscle cells.

25 Preferably, the reduction or delay in apoptosis would be at least a 10% reduction or delay, including, for example, 15%, 20%, 25%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100% or any amount in between. The reduction or delay can be measured, for example, by comparing the number of cells after contact with the tetrahydrobenzothiazole analogue to the number of cells in a control population of cells lacking contact with the tetrahydrobenzothiazole analogue. Histological signs of apoptosis that would be reduced or delayed in cells after contact with the tetrahydrobenzothiazole analogue include condensation of the chromatin, the occurrence of apoptotic bodies, and cellular shrinkage. DNA laddering and other signs of DNA degradation are also signs of apoptosis which would be reduced or delayed after contact with the tetrahydrobenzothiazole analogue. Apoptosis can also be assessed indirectly by observing, for example, a reduction in the amount of release or

5 activity by the population of cells. Thus, if the cell population undergoes apoptosis, neurotransmitter release upon stimulation of neuronal cells would decrease. Similarly, a decrease in cardiac muscle contraction or cardiac output is an indicator of apoptosis. It is understood that one or a combination of indicators of apoptosis may show a delay or reduction.

10

The cell can be contacted *in vitro* with the agent, for example, by adding the agent to the culture medium (by continuous infusion, by bolus delivery, or by changing the medium to a medium that contains the agent) or by adding the agent to the extracellular fluid *in vivo* (by local delivery, systemic delivery, intravenous injection, 15 bolus delivery, or continuous infusion). *In vitro* contact may be preferred, for example, in reducing or delaying apoptosis in a population of cells to be transplanted into a donor. *In vivo* contact may be preferred in reducing or delaying apoptosis in a subject with a disease, condition, or injury associated with apoptosis. The duration of “contact” with a cell or population of cells is determined by the time the agent is 20 present at physiologically effective levels or at presumed physiologically effective levels in the medium or extracellular fluid bathing the cell or cells. Preferably, the duration of contact is 1-96 hours and, more preferably, for 24 hours, but such time would vary based on the half life of the agent and could be optimized by one skilled in the art using routine experimentation.

25

It is understood that the tetrahydrobenzothiazole analogue is administered in a therapeutically effective dose or doses to reduce or delay apoptosis. The agents used in this invention (i.e., tetrahydrobenzothiazole analogues) can be administered *in vitro* in an amount of about 10 nmol to 100µmol. More preferably the agent is administered *in* 30 *vitro* in an amount of about 1.0 nmol to 10µmol. Necessary modifications in this dosage range may be determined by one of ordinary skill in the art.

The agents used in this invention are administered *in vivo* to a subject in need thereof by commonly employed methods for administering agents in such a way to 35 bring the agent in contact with the population of cells. The agents of the present

5 invention can be administered orally, parenterally, transdermally, extracorporeally, topically or the like, although intravenous administration is typically preferred.

The agents of the present invention can also be administered using gene therapy methods of delivery. See, e.g., U.S. Patent No. 5,399,346, which is incorporated by
10 reference herein. Using a gene therapy method of delivery, primary cells transfected with the gene for the agent of the present invention can additionally be transfected with tissue specific promoters to target specific organs, tissue, grafts, or cells.

The dosage of the agent varies depending on the type of disease or condition,
15 degree of apoptosis, weight, age, sex, and method of administration. Also, the dosage of the agent varies depending on the target cell, tissue, graft, or organ. Generally, the agents can be orally or intravenously administered *in vivo* in an amount of about 0.01-1000 mg/kg. More preferably, the agent is administered *in vivo* in an amount of about 0.1 to 100 mg/kg. Even more preferably, the agent is administered in an amount of
20 about 5mg/kg. Thus, an administration regimen could include long-term, daily treatment. By "long-term" is meant at least two weeks and, preferably, several weeks, months, or years of duration. Necessary modifications in this dosage range may be determined by one of ordinary skill in the art using only routine experimentation given the teachings herein. See *Remington's Pharmaceutical Sciences* (Martin, E.W., ed.,
25 latest edition), Mack Publishing Co., Easton, PA. The dosage can also be adjusted by the individual physician in the event of any complication.

The agents can be administered conventionally as compositions containing the active agent as a predetermined quantity of active material calculated to produce the
30 desired therapeutic effect in association with the required diluent, i.e., carrier or vehicle. Depending on the intended mode of administration, the agent can be in pharmaceutical compositions in the form of solid, semi-solid or liquid dosage forms, such as, for example, tablets, suppositories, pills, capsules, powders, liquids, suspensions, lotions, creams, gels, or the like, preferably in unit dosage form suitable
35 for single administration of a precise dosage. The compositions of Formulas (I), (II), (III), (IV), (V), (VI), (VII), (VIII), (IX), (X) (XI) through (XV) include, as noted above,

- 5 an effective amount of the selected compound in combination with a pharmaceutically acceptable carrier and, in addition, may include other medicinal agents, pharmaceutical agents, carriers, adjuvants, diluents, etc.

For solid compositions, conventional nontoxic solid carriers include, for
10 example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharin, talc, cellulose, glucose, sucrose, magnesium carbonate, and the like. Liquid pharmaceutically administrable compositions can, for example, be prepared by dissolving, dispersing, etc. an active compound as described herein and optional pharmaceutical adjuvants in an excipient, such as, for example, water, saline, aqueous
15 dextrose, glycerol, ethanol, and the like, to thereby form a solution or suspension. If desired, the pharmaceutical composition to be administered may also contain minor amounts of nontoxic auxiliary substances such as wetting or emulsifying agents, pH buffering agents and the like, for example, sodium acetate, sorbitan monolaurate, triethanolamine sodium acetate, triethanolamine oleate, etc. Thus, the compositions are
20 administered in a manner compatible with the dosage formulation and in a therapeutically effective amount. As discussed above, precise amounts of active ingredient required to be administered depend on the judgment of the practitioner and are peculiar to each individual.

25 For oral administration, fine powders or granules may contain diluting, dispersing, and/or surface active agents, and may be presented in water or in a syrup, in capsules or sachets in the dry state, or in a nonaqueous solution or suspension wherein suspending agents may be included, in tablets wherein binders and lubricants may be included, or in a suspension in water or a syrup. Where desirable or necessary,
30 flavoring, preserving, suspending, thickening, or emulsifying agents may be included. Tablets and granules are preferred oral administration forms, and these may be coated.

Parenteral administration, if used, is generally characterized by intradermal, subcutaneous, intramuscular, intraperitoneal, intravenous, intra-articular and
35 intratracheal routes of injection. A more recently revised approach for parenteral administration involves use of a slow release or sustained release system such that a

5 constant dosage is maintained. See, e.g., U.S. Patent No. 3,610,795, which is
incorporated by reference herein. The agents can also be administered using polymer
based delivery systems, including, for example, microencapsulation as described in
Langer (1998). Injectables can be prepared in conventional forms, either as liquid
10 solutions or suspensions, solid forms suitable for solution or suspension in liquid prior
to injection, or as emulsions.

For topical administration, liquids, suspension, lotions, creams, gels or the like
may be used as long as the active compound can be delivered to the surface of the skin.

15 Experimental

The following examples are put forth so as to provide those of ordinary skill in
the art with a complete disclosure and description of how the compounds,
compositions, articles, devices, and/or methods claimed herein are made and evaluated,
and are intended to be purely exemplary of the invention and are not intended to limit
20 the scope of what the inventors regard as their invention. Efforts have been made to
ensure accuracy with respect to numbers (e.g., amounts, temperature, etc.) but some
errors and deviations should be accounted for. Unless indicated otherwise, parts are
parts by weight, temperature is in °C or is at ambient temperature, and pressure is at or
near atmospheric.

25

The dotted lines attached to the benzene ring of structures 4-25 in the examples
denote a chemical bond and not an additional carbon atom. Thus the R group of
structure 4 is chemically bonded to Formula (XI) at the *para* position relative to the
Fluorine atom. Throughout these examples, the dotted lines are meant to be a bond
30 attachment and not an additional carbon atom.

Numbers in bold and parenthesis correspond to structures depicted in the
examples for formula (XI) of example 1 and formula (XII) of example 2.

5

Synthesis of the intermediate 2-amino-4,5,6,7-tetrahydrobenzothiazole

A mixture of cyclohexanone (1.96 g, 0.02 mmol), thiourea (3.04 g, 0.04 mmol) and iodine (5.08 g, 0.02 mmol) was stirred in a 110°C oil bath for 12 h. The reaction mixture then was cooled, dissolved in boiling water and extracted with ether to remove any ketone and iodine. The solution was made basic with solid NaHCO₃, yellow crystals thereafter precipitated and were collected by filter to give 2-amino-4,5,6,7-tetrahydrobenzothiazole hydrogen iodide (3.2 g, 57%), m.p. 185-187°C. ¹H NMR (DMSO-d₆): 2.50-2.48 (m, 4 H), 1.78-1.75 (m, 4 H). ¹³C NMR (DMSO-d₆): 168.6, 140.7, 116.3, 26.1, 24.1, 24.0, 23.6. 2-amino-4,5,6,7-tetrahydrobenzothiazole hydrogen iodide (0.5 g) then was dissolved in hot saturated aqueous Na₂CO₃, and on cooling 2-amino-4,5,6,7-tetrahydrobenzothiazole was gained as white needle crystals (0.25g, 93%), m.p. 87-88°C.

20

Example 1

Synthesis of Z-1-110: 1-(4,5,6,7-tetrahydro-2-imino-3(2H)-benzothiazolyl)-2-butanone hydrobromide (1)

A mixture of 2-amino-4,5,6,7-tetrahydrobenzothiazole (76 mg, 0.49 mmol) and 1-bromo-2-butanone (51 µl, 0.50 mmol) in benzene (10 ml) was stirred at room temperature for two days. The resulting precipitate was filtered off from the reaction mixture and washed with a small amount of ethanol and benzene to give 1-(4,5,6,7-tetrahydro-benzothiazolyl)-2-butanone hydrobromide (1) (75 mg, 50%) as pale yellow crystals. mp 114-115°C ¹H NMR (DMSO-d₆): δ 9.45 (s, 1 H), 5.06 (s, 2 H), 2.64 (q, J = 7.2 Hz, 2 H), 2.41-2.52 (m, 2 H), 2.29 (br, 2H), 1.73 (br, 4H), 0.99 (t, J = 7.2 Hz, 3 H); Anal. Calcd. HR-MS m/z calcd for C₁₁H₁₇N₂OS 225.1062, found 25.1071. Anal. Calcd. C₁₁H₁₇BrN₂OS: C, 43.18; H, 5.61. Found: C, 43.23; H, 5.56.

5 Synthesis of PFT- α (Z-1-073): 1-(4-methylphenyl)-2-(4,5,6,7-tetrahydro-2-imino-3(2H)-benzothiazolyl)-ethanone hydrobromide (5)

A mixture of 2-amino-4,5,6,7-tetrahydrobenzothiazole (2) (154 mg, 1 mmol) and 2-bromo-4'-methylacetophone (213 mg, 1 mmol) in dry benzene (20 ml) was stirred at
10 room temperature for two days. Thereafter, the precipitate was filtered off from the reaction mixture and washed with a small amount of benzene to afford 1-(4-methylphenyl)-2-(4,5,6,7-tetrahydro-2-imino-3(2H)-benzothiazolyl)-ethanone hydrobromide (5) (247 mg, 67%) as white crystals. mp 180°C (EtOH/EtOAc) (lit²⁰: 182°C); ¹H NMR (DMSO-d₆): δ 8.85 (s, 1 H), 7.96 (d, J = 8.1 Hz, 2 H), 7.45 (d, J = 8.1
15 Hz, 2 H), 5.70 (s, 2 H), 2.55-2.30 (m, 7 H), 1.73 (m, 4 H).

The following compounds were synthesized in accordance with the methods above, except the products were crystallized from MeOH/EtOAc, Z-1-189: ethyl 2-(4,5,6,7-tetrahydro-2-imino-3(2H)-benzothiazolyl)-acetate hydrobromide (2),: yield (52%); mp
20 224°C (MeOH/EtOAc); ¹H NMR (DMSO-d₆) δ 9.63 (s, 1H), 4.93 (s, 2H), 4.21 (q, J = 7.1Hz, 2H), 2.38 (m, 2H), 1.73 (m, 4H), 1.24 (t, J = 7.1Hz, 3H),

ethyl 2-(4,5,6,7-tetrahydro-2-imino-3(2H)-benzothiazolyl)-acetate hydrohalide (2a),
25 methyl 2-(4,5,6,7-tetrahydro-2-imino-3(2H)-benzothiazolyl)-acetate hydrohalide (3),

Z-1-135: 1-(4-fluorophenyl)-2-(4,5,6,7-tetrahydro-2-imino-3(2H)-benzothiazolyl)-ethanone hydrobromide (4): 44%, ¹H NMR (DMSO-d₆): 9.51 (s, 1 H), 8.14 (d, J = 5.5 Hz and J = 8.7 Hz, 2 H), 7.49 (t, J = 8.7 Hz, 2 H), 5.74 (s, 2 H),
30 2.55-2.34 (m, 4 H), 1.73 (m, 4 H). HR-MS m/z calcd for C₁₅H₁₆FN₂OS 291.0967, found 291.0964. Anal. (C₁₅H₁₆BrFN₂OS) C, H, N,

1-(4-fluorophenyl)-2-(4,5,6,7-tetrahydro-2-imino-3(2H)-benzothiazolyl)-ethanone hydrohalide (4a),
35

- 5 Z-1-117: 1-(4-methylphenyl)-2-(4,5,6,7-tetrahydro-2-imino-6-methyl-3(2*H*)-benzothiazolyl)-ethanone hydrobromide (5x): 54%; mp 254-256°C (lit²⁰ 278°C); ¹H NMR (DMSO-*d*₆): 9.60 (s, 1 H), 7.95 (d, *J* = 8.2 Hz, 2 H), 7.45 (d, *J* = 8.2 Hz, 2 H), 5.70 (s, 2 H), 2.71-2.64 (m, 1 H), 2.51-2.31 (m, 5 H), 2.21-2.12 (m, 1 H), 1.83-1.79 (m, 2 H), 1.36 (br, 1 H), 1.02 (d, *J* = 6.5 Hz, 3 H). Anal. (C₁₇H₂₁BrN₂OS) C, H, N,
- 10 1-(4-methylphenyl)-2-(4,5,6,7-tetrahydro-2-imino-3(2*H*)-benzothiazolyl)-acetate hydrohalide (5xx),
- Z-1-133: 1-(4-methoxyphenyl)-2-(4,5,6,7-tetrahydro-2-imino-3(2*H*)-benzothiazolyl)-ethanone hydrobromide (6): 61%; ¹H NMR (DMSO-*d*₆): 9.47 (s, 1 H), 8.03 (d, *J* = 8.9 Hz, 2 H), 7.16 (d, *J* = 8.9 Hz, 2 H), 5.67 (s, 2 H), 3.89 (s, 3 H), 2.55 (br, 2 H), 2.32 (br, 2 H), 1.73 (br, 4 H). HR-MS *m/z* calcd for C₁₆H₁₉N₂O₂S 303.1167, found 303.1158. Anal. (C₁₆H₁₉BrN₂O₂S) C, H, N,
- 15 1-(4-methoxyphenyl)-2-(4,5,6,7-tetrahydro-2-imino-3(2*H*)-benzothiazolyl)-ethanone hydrohalide (6a),
- Z-1-138: 1-(2-methoxyphenyl)-2-(4,5,6,7-tetrahydro-2-imino-3(2*H*)-benzothiazolyl)-ethanone hydrobromide (7): 17%, ¹H NMR (DMSO-*d*₆): 7.70-7.12 (m, 4 H), 5.42 (s, 2 H), 3.99 (s, 3 H), 2.50-2.39 (m, 4 H), 1.81 (br, 4 H). HR-MS *m/z* calcd for C₁₆H₁₉N₂O₂S 303.1167, found 303.1173. Anal. (C₁₆H₁₉BrN₂O₂S) C, H, N,
- 25 1-(2-methoxyphenyl)-2-(4,5,6,7-tetrahydro-2-imino-3(2*H*)-benzothiazolyl)-ethanone hydrohalide (7a),
- 30 Z-1-139: 1-(3-methoxyphenyl)-2-(4,5,6,7-tetrahydro-2-imino-3(2*H*)-benzothiazolyl)-ethanone hydrobromide (8): 52%, mp 143-145°C; ¹H NMR (DMSO-*d*₆): 9.50 (s, 1 H), 7.67-7.37 (m, 4 H), 5.74 (s, 2 H), 3.86 (s, 3 H), 2.55-2.33 (m, 4H), 1.73 (br, 4 H). HR-MS *m/z* calcd for C₁₆H₁₉N₂O₂S 303.1167, found 303.1171. Anal. (C₁₆H₁₉BrN₂O₂S) C,
- 35 H, N,

- 5 1-(3-methoxyphenyl)-2-(4,5,6,7-tetrahydro-2-imino-3(2*H*)-benzothiazolyl)-ethanone hydrohalide (8a),

Z-1-141: 1-(4-chlorophenyl)-2-(4,5,6,7-tetrahydro-2-imino-3(2*H*)-benzothiazolyl)-ethanone hydrobromide (9): 70%; ¹H NMR (DMSO-*d*₆): 9.52 (s, 1 H), 3.06 (d, *J* = 8.6 Hz, 2 H), 7.73 (d, *J* = 8.6 Hz, 2 H), 5.74 (s, 2 H), 2.55-2.33 (m, 4 H), 1.72 (br, 4 H),

1-(4-chlorophenyl)-2-(4,5,6,7-tetrahydro-2-imino-3(2*H*)-benzothiazolyl)-acetate hydrohalide (9a),

- 15 Z-1-145: 1-(4-phenylphenyl)-2-(4,5,6,7-tetrahydro-2-imino-3(2*H*)-benzothiazolyl)-ethanone hydrobromide (10) : 58%; ¹H NMR (DMSO-*d*₆): 9.55 (s, 1 H), 8.15(d, *J* = 8.5 Hz, 2 H), 7.81 (d, *J* = 7.0 Hz, 2 H), 7.57-7.44 (m, 4 H), 5.80 (s, 2 H), 2.56-2.36 (m, 4 H), 1.74 (br, 4 H),

- 20 1-(4-phenylphenyl)-2-(4,5,6,7-tetrahydro-2-imino-3(2*H*)-benzothiazolyl)-acetate hydrohalide (10a),

Z-1-153: 1-(4-nitrophenyl)-2-(4,5,6,7-tetrahydro-2-imino-3(2*H*)-benzothiazolyl)-ethanone hydrobromide (11): 61%, ¹H NMR (DMSO-*d*₆): 9.54 (s, 1 H), 8.46 (d, *J* = 8.9 Hz, 2 H), 8.28 (d, *J* = 8.9 Hz, 2 H), 5.80 (s, 2 H), 2.56-2.37 (m, 4 H), 1.74 (br, 4 H).
25 HR-MS *m/z* calcd for C₁₅H₁₆N₃O₃S 318.0912, found 318.0903. Anal. (C₁₅H₁₆BrN₃O₃S) C, H, N,


- 30 1-(4-nitrophenyl)-2-(4,5,6,7-tetrahydro-2-imino-3(2*H*)-benzothiazolyl)-ethanone hydrohalide (11a),

1-(4-tetradecyloxyphenyl)-2-(4,5,6,7-tetrahydro-2-imino-3(2*H*)-benzothiazolyl)-ethanone hydrohalide (12),

- 35 Z-1-143: 1-phenyl-2-(4,5,6,7-tetrahydro-2-imino-3(2*H*)-benzothiazolyl)-ethanone hydrobromide (13): 58%, mp 151-152°C; ¹H NMR (DMSO-*d*₆): 9.53 (s, 1 H), 8.07-

- 5 8.04 (m, 2 H), 7.79-7.74 (m, 1 H), 7.66-7.62 (m, 2 H), 5.75 (s, 2 H), 2.55-2.33 (m, 4H), 1.72 (br, 4 H). HR-MS m/z calcd for $C_{15}H_{17}N_2OS$ 273.1162, found 273.1058. Anal. ($C_{15}H_{17}BrN_2OS$) C, H, N,

- 10 1-phenyl-2-(4,5,6,7-tetrahydro-2-imino-3(2*H*)-benzothiazolyl)-ethanone hydrohalide (13a), and



Example 2

- 15 Synthesis of Z-1-119: 3-(phenylmethyl)-4,5,6,7-tetrahydro-2(3*H*)-benzothiazolimine hydrobromide (14)

5 A mixture of 2-amino-4,5,6,7-tetrahydrobenzothiazole (101 mg, 0.66 mmol) and benzyl bromide (115 mg, 0.67 mmol) in THF (5 mL) was refluxed for two days. The precipitate was collected by filter, washed with a small amount of THF, and recrystallized from ethanol-ethyl ether to afford compound (14) (75 mg, 75%) as pale yellow crystals; mp 271°C; ¹H NMR (DMSO-d₆): 9.63 (s, 1 H), 7.45-7.35 (m, 3 H),
10 7.15 (d, J = 7.1 Hz, 2 H), 5.29 (s, 2 H), 2.50-2.30 (m, 4 H), 1.76 (br, 4 H); HR-MS m/z calcd for C₁₄H₁₇N₂S 245.1112, found 245.1104. Anal. (C₁₄H₁₇BrN₂S) C, H, N.

2-imino-3-benzyl-4,5,6,7-tetrahydro-3(2H)- benzothiazole hydrohalide (14a).

15 Synthesis of Z-1-113: 3-(cyclopropylmethyl)-4,5,6,7-tetrahydro-2(3H)-benzothiazolimine hydrobromide (15)

A mixture of 2-amino-4,5,6,7-tetrahydrobenzothiazole (52 mg, 0.34 mmol) and (bromomethyl)cyclopropane (47 mg, 97%, 0.34 mmol) in ethanol (5 mL) was refluxed
20 for one day. Thereafter, the solvent was evaporated under vacuum. The residue was extracted with hot benzene, filtered and washed with benzene to gain a pale yellow solid that then was recrystallized from ethanol-ethyl ether to afford (15) (60 mg, 61%) as pale yellow crystals; mp 239-240°C; ¹H NMR (DMSO-d₆): 3.88 (d, J = 7.0 Hz, 2 H), 2.52-2.39 (m, 4 H), 1.76-1.71 (m, 4 H), 1.09 (br, 1 H), 0.55-0.53 (m, 2 H), 0.42-0.40
25 (m, 2 H); HR-MS m/z calcd for C₁₁H₁₇N₂S 209.1112, found 209.1115. Anal. (C₁₁H₁₇BrN₂S) C, H, N.

2-imino-3-cyclopropylmethyl-4,5,6,7-tetrahydro-3(2H)-benzothiazole hydrohalide (15a),

30

The following compounds were synthesized in accordance with the methods for the synthesis of compounds (19) Z-1-165.

Z-1-165: 2-imino-3-(4'-tert-butylbenzyl)-4,5,6,7-tetrahydro-3(2H)-benzothiazole hydrobromide (19)

35

5 A mixture of 2-amino-4,5,6,7-tetrahydrobenzothiazole (101 mg, 0.66 mmol) and 4-(*tert*-butyl) benzyl bromide (154 mg, 0.66 mmol) in THF (5 ml) was refluxed for two days. The resulting precipitate was collected by filter and washed with a small amount of THF and EtOH, and recrystallized from MeOH/EtOAc to give 2-imino-3-(4-*tert*-butylbenzyl)-4,5,6,7-tetrahydro-3(2H)-benzothiazole hydrobromide (98 mg, 39%)
10 as white crystals.

mp 255-257°C (MeOH/EtOAc); ¹H NMR (DMSO-d₆) δ 9.55 (s, 1H), 7.43 (d, J = 8.4 Hz, 2H), 7.07 (d, J = 8.4 Hz, 2H), 5.22 (s, 2H), 2.38 (m, 2H), 1.72 (m, 4H), 1.26 (s, 9H); Anal. (C₁₈H₂₅BrN₂S·0.25H₂O) C, H, N,

15

Z-1-167: 2-imino-3-phenylethyl-4,5,6,7-tetrahydro-3(2H)-benzothiazole (25),

The mixture of 2-amino-4,5,6,7-tetrahydrobenzothiazole (110 mg), 2-bromoethylbenzene (135 mg), KI (110 mg) in ethanol (5 mL) was refluxed for one day.
20 The solvent was evaporated under vacuum. The residue was neutralized with saturated aqueous Na₂CO₃ solution. The product was separated from the aqueous solution as an uncrystallizable oil, and recrystallized from MeOH/EtOAc to give compound as pale brown crystal (yield 38%); mp 262°C (MeOH/EtOAc); ¹H NMR (DMSO-d₆) δ 9.33 (br, 1H), 7.32-7.20 (m, 5H), 4.06 (t, J = 6.9 Hz, 2H), 2.92 (t, J = 6.9 Hz, 2H), 2.42 (m,
25 2H), 2.14 (m, 2H), 1.61 (m, 4H),

2-imino-3-phenylethyl-4,5,6,7-tetrahydro-3(2H)-benzothiazole hydrohalide (25a),

Z-1-181 2-imino-3-(4'-fluorobenzyl)-4,5,6,7-tetrahydro-3(2H)-benzothiazole
30 hydrobromide (16), yield (41%); mp 252°C (MeOH/EtOAc); ¹H NMR (DMSO-d₆) δ 9.61 (s, 1H), 7.30-7.23 (m, 4H), 5.26 (s, 2H), 2.36 (m, 2H), 1.71 (m, 4H),

2-imino-3-(4-fluorobenzyl)-4,5,6,7-tetrahydro-3(2H)-benzothiazole hydrohalide (16a),

35 Z-1-163: 2-imino-3-(2',4'-difluorobenzyl)-4,5,6,7-tetrahydro-3(2H)-benzothiazole hydrobromide (17),: yield (45%) ; mp 263-264°C (MeOH/EtOAc); ¹H NMR (DMSO-

5 d₆) δ 9.66 (s, 1H), 7.42-7.36 (m, 1H), 7.15-7.11 (m, 2 H), 5.28 (s, 2H), 2.33 (m, 2H), 1.70 (m, 4H),

2-imino-3-(2,4-difluorobenzyl)-4,5,6,7-tetrahydro-3(2*H*)-benzothiazole hydrohalide (17a),

10

Z-1-179: 2-imino-3-(4'-methylbenzyl)-4,5,6,7-tetrahydro-3(2*H*)-benzothiazole hydrobromide (18), : yield (39%) ; mp 253°C (MeOH/EtOAc); ¹H NMR (DMSO-d₆) δ 9.58 (s, 1H), 7.23 (d, J = 7.9 Hz, 2H), 7.05 (d, J = 7.9 Hz, 2H), 5.22 (s, 2H), 2.35 (m, 2H), 2.29 (s, 3H), 1.70 (m, 4H),

15

2-imino-3-(4-methylbenzyl)-4,5,6,7-tetrahydro-3(2*H*)-benzothiazole hydrohalide (18a),

Z-1-161: 2-imino-3-(4'-nitrobenzyl)-4,5,6,7-tetrahydro-3(2*H*)-benzothiazole hydrobromide (20),: yield (44%); mp 235-236°C (MeOH/EtOAc); ¹H NMR (DMSO-d₆) δ 9.75 (s, 1H), 8.37 (d, J = 8.8Hz, 2H), 7.51 (d, J = 8.8 Hz, 2H), 5.52 (s, 2H), 2.59 (m, 2H), 2.41 (m, 2H), 1.80 (m, 4H),

20

2-imino-3-(4-nitrobenzyl)-4,5,6,7-tetrahydro-3(2*H*)-benzothiazole hydrohalide (20a),

25 Z-1-183: 2-imino-3-(4'-cyanobenzyl)-4,5,6,7-tetrahydro-3(2*H*)-benzothiazole hydrobromide (21), : yield (32%); mp 247°C (MeOH/EtOAc); ¹H NMR (DMSO-d₆) δ 9.64 (s, 1H), 7.94 (d, J = 8.3 Hz, 2H), 7.35 (d, J = 8.3 Hz, 2H), 5.38 (s, 2H), 2.33 (m, 2H), 1.73 (m, 4H),

30 2-imino-3-(4-cynobenzyl)-4,5,6,7-tetrahydro-3(2*H*)-benzothiazole hydrohalide (21a),

Z-1-175: 2-imino-3-(4'-trifluorobenzyl)-4,5,6,7-tetrahydro-3(2*H*)-benzothiazole hydrobromide (22), : yield (40%); mp 265-266°C (MeOH/EtOAc); ¹H NMR (DMSO-d₆) δ 9.68 (s, 1H), 7.80 (d, J = 8.0Hz, 2H), 7.38 (d, J = 8.0Hz, 2H), 5.41 (s, 2H), 2.55 (m, 2H), 2.35 (m, 2H), 1.72 (m, 4H); Anal. (C₁₅H₁₆BrF₃N₂S·H₂O) C, H, N,

35

5

2-imino-3-(4-trifluoromethylbenzyl)-4,5,6,7-tetrahydro-3(2*H*)-benzothiazole hydrohalide (22a),

10

2-imino-3-(4-phenylbenzyl)-4,5,6,7-tetrahydro-3(2*H*)-benzothiazole hydrohalide (23),
Z-1-171: 2-imino-3-(naphthalenylmethyl)-4,5,6,7-tetrahydro-3(2*H*)-benzothiazole hydrobromide (24), : yield (34%); mp 268°C (MeOH/EtOAc); ¹H NMR (DMSO-d₆) δ 9.66 (s, 1H), 8.0-7.92 (m, 3H), 7.64 (s, 1H), 7.56-7.53 (m, 2H), 7.33 (d, J = 7.2 Hz, 1H), 5.44 (s, 1H), 2.54 (m, 2H), 2.39 (m, 2H), 1.68 (m, 4H).

15

2-imino-3-(2-menaphthyl)-4,5,6,7-tetrahydro-3(2*H*)-benzothiazole hydrohalide (24a),

Z-1-187: 2-Imino-3-butyl-4,5,6,7-tetrahydro-3(2*H*)-benzothiazole hydrobromide (26), yield (19%); ¹H NMR (DMSO-d₆) δ 9.34 (s, 1H), 3.87 (t, J = 7.1 Hz, 2H), 1.77 (m, 4H), 1.57 (m, 2H), 1.34 (m, 2H), 0.91 (t, J = 7.1 Hz, 3H),

20














2-imino-3-butyl-4,5,6,7-tetrahydro-3(2*H*)-benzothiazole hydrohalide (26a),

Z-2-007: 1-(4-methylphenyl)-2-(4,5,6,7-tetrahydro-2-imino-3(2*H*)-benzoxazolyl)-ethanone hydrobromide (27): yield (51%); mp 237°C (MeOH/EtOAc); ¹H NMR (DMSO-d₆) δ 9.64 (br, 1H), 7.95 (d, J = 8.0Hz, 2H), 7.46 (d, J = 8.9Hz, 2H), 5.78 (s, 2H), 2.52 (br, 2H), 2.44 (s, 3H), 2.33 (m, 2H), 1.77 (m, 4H)

25


Z-2-013: 1-(4,5,6,7-tetrahydro-2-imino-3(2*H*)-benzoxazolyl)-2-butanone hydrobromide (28): yield (57%); mp 206°C (MeOH/EtOAc); ¹H NMR (DMSO-d₆) δ 9.56 (s, 1H), 4.95 (s, 2H), 2.59 (q, J = 7.2Hz, 2H), 2.27 (m, 2H), 1.78-1.69 (m, 4H), 0.99 (t, J = 7.2 Hz, 3H),

30



5

Scheme A



10

Z-2-035II: 2-(4'-methylphenyl)-5,6,7,8-tetrahydroimidazo[2,1-b]benzothiazole (29), mp 187°C (MeOH/EtOAc) (lit 185°C); ¹H NMR (DMSO-d₆) δ 7.71 (d, J = 8.1Hz, 2H), 7.50 (s, 1H), 7.19 (d, J = 8.1Hz, 2H), 2.70-2.63 (m, 4H), 2.35 (s, 3H), 1.95-1.93 (m, 4H).

15

Compound	R ₁	X
29	H CH ₃	S O

wherein R₂ is C₁₋₃ alkyl or aryl substituted or unsubstituted


20

Example 3

A mixture of 2, 3, 4, 5, 6, 7-hexahydrobenzothiazolin-2-one (0.1 mol) and sodium hydride (0.1 mol) in toluene (20 ml) was refluxed for half an hour. 2-bromo-4'-methylacetophenone (0.1 mol) was added and the reaction mixture was refluxed for an additional four hours. The precipitated sodium bromide was filtered off and the toluene was removed under vacuum. The crude product was purified to give 3-*p*-methylphenacyl-2, 3, 4, 5, 6, 7-hexahydrobenzothiazolin-2-one (XIII) in the yield of 48%. HNMR (CDCl₃): 7.88 (d, J=8.0 Hz, 2 H), 7.28 (d, J=8.0 Hz, 2 H), 5.04 (s, 2 H), 2.42 (br, 5 H), 2.19-2.17 (m, 2 H), 1.82-1.78 (m, 4 H).

25

5



(XIII)

10

Example 4**Neuroprotective Activity of PFT α , its Precursor, and Analogues in PC12 cells**

The 2-amino-4,5,6,7-tetrahydrobenzothiazole hydrogen iodide, 1-(4-methylphenyl)-2-(4,5,6,7-tetrahydro-2-imino-3(2*H*)-benzothiazolyl)-ethanone hydrobromide, and analogues were tested for their ability to protect PC12 cells against the toxic agent, camptothecin. See Figures 2a and 2b. PC12 cells were cultured in RPMI1640 media containing 10% horse serum and 5% fetal calf serum (37°C, pH 7.4) until confluent. Thereafter, 2.2×10^4 cells per mL were added to each trough of a 96 well plate (approximately 6,270 cells per well 285 μ L) and were left to grow for 12 h.

The media then was removed and was replaced by media with and without the compounds to be tested, including precursor, PFT α , and numerous analogues (concentration 100-400 nM, prepared in DMSO to a final dilution of 0.4%, 8 wells per concentration). Following 6 h of incubation, camptothecin (40 μ M) was added to 4 of each 8 wells, and the cells were incubated for a further 24 h. Cells then were gently washed twice with phosphate buffered saline (PBS: 0.1 M, 37°C, pH 7.4) and were incubated in PBS containing the live-cell indicator dye, 2', 7'-bis(2-carboxyethyl)-5-6-carboxyfluorescein AM ester (BCECF AM: 5 μ M) (Molecular Probes, CA), for 45 min in minimal light. This fluorescent dye was taken up and retained by live cells. Finally,

5 the cells were washed twice in PBS and their fluorescence then was quantified (excitation: 490nm, emission: 535nm, Perkin Elmer HTS 7000). See Figure 2A.

PC12 cell survival was also assessed. The results are shown in Figure 2B. One hundred percent cell survival was determined from cells incubated in the absence of both any compound and camptothecin. One hundred percent cellular death was
10 determined from cells incubated in the absence of any compound and presence of camptothecin, which in prior studies was found to occur at a minimum concentration of 40 μ M. Compound-induced toxicity was assessed by comparing cells treated with compound at various concentrations in the absence of camptothecin with cells incubated in the absence of both compound and camptothecin. In the presence of PFT α ,
15 neuronal survival of PC12 cells was increased up to 70% as compared to the PC12 cells treated with the DNA damaging agent, camptothecin, in the absence of PFT α . The precursor, 2-amino-4,5,6,7-tetrahydrobenzothiazole hydrogen iodide, showed low activity under the same conditions. No compound was found to induce PC12 cell toxicity at the maximal concentration assessed (400 nM). Compound-induced cellular
20 protection was assessed by comparing cells incubated with camptothecin with cells incubated with camptothecin in the presence of compound, in a concentration-dependent manner. Finally, the natural fluorescence of each compound was quantified in the absence of cells and BCECF AM, and was found to be negligible. A 50% protective concentration (PC₅₀), commensurate with an IC₅₀ (concentration required to
25 inhibit 50% of p53 activity), was calculated by transforming the data into a logit format (logit = $\ln(\% \text{ protection}/100\% - \% \text{ protection})$), and calculating the value from a correlation between a plot of the log concentration of compound versus logit activity.

In summary, compounds 5x, 6-7, 9, 13, proved to be highly potent in protecting PC12 cells. The concentration of each analogue required to protect 50% of cells
30 (determined as a PC₅₀ value) is shown in Table 1.

5

Table 1

Concentration of compounds required to induce 50% protection (PC₅₀) of PC₁₂ cells from camptothecin-induced cell-death.

10	Compound		PC ₅₀ nM
	(precursor)		5754
	5	(PFTα)	252
	14	(Z-119)	741
	1	(Z-110)	571
15	15	(Z-113)	728
	5x	(Z-117)	306
	6	(Z-133)	182
	7	(Z-138)	348
	8	(Z-139)	418
20	13	(Z-143)	169
	10	(Z-145)	767
	4	(Z-135)	379
	9	(Z-141)	187
	11	(Z-153)	410
25	16	(Z-1-181)	154
	17	(Z-1-163)	93
	18	(Z-1-179)	1000
	19	(Z-1-165)	315
	20	(Z-1-161)	144
30	21	(Z-1-183)	418
	22	(Z-1-175)	327
	24	(Z-1-171)	318
	25	(Z-1-167)	1100
	26	(Z-1-187)	395
35	2	(Z-1-189)	595
	30	(Z-1-169)	133

5	27	(Z-2-007)	69
	28	(Z-2-013)	143
	29	(Z-2-035II)	214

In summary, numerous of the analogues (e.g. 5x, 6-7, 9-13, etc.) prove to be
10 highly potent.

Example 5

Neuroprotective Activity of PFT α and its Precursor in Rat Hippocampal Neurons

In Vitro

15

2-amino-4,5,6,7-tetrahydrobenzothiazole hydrogen and 1-(4-methylphenyl)-2-
(4,5,6,7-tetrahydro-2-imino-3(2*H*)-benzothiazolyl)-ethanone hydrobromide and their
analogues were assessed for their ability to protect rat hippocampal cells in culture
20 from camptothecin (5 μ M) or etoposide (5 μ M). See Figures 3-5.

Dissociated hippocampal cell cultures were established from embryonic day 18
Sprague-Dawley rats (Harlan, Inc.) as described previously. See Culmsee et al. (2001)
J. Neurochem. 77:220-28. Cells were grown in polyethyleneimine-coated plastic
dishes or 22 mm² glass coverslips, and incubated in Neurobasal medium containing B-
25 27 supplements, 2 mM L-glutamine, 25 mg/ml gentamycin, 1 mM Hepes (Gibco BRL)
and 0.001% gentamicin sulfate. All experiments were performed using 9-10 day-old
cultures. Camptothecin and etoposide (Sigma) were prepared as 500X stocks in DMSO.
Neuron survival was quantified by established methods. See Mattson et al. (1993)
Neuron 10:243-54. Briefly, viable neurons in premarked fields (10X objective) were
30 counted before experimental treatment and at specified time points thereafter. Neurons
with intact neurites of uniform diameter and soma with a smooth round appearance
were considered viable. In contrast neurons with fragmented neurites and vacuolated
soma were considered non-viable.

The results showed that PFT α (5) possessed potent activity to protect the primary
35 hippocampal cells. At concentrations up to 400 nM, neither PFT α (5) nor its
precursor were toxic when administered alone to cells. However, both camptothecin (5

5 μM) and etoposide ($2.5 \mu\text{M}$), known to induce neuronal apoptosis by a mechanism involving DNA damage and p53 induction, killed approximately 85% of neurons. Pretreatment of cells with PFT α (5), but not its precursor, resulted in improved cell survival compared to vehicle controls.

10 The neuroprotective effect of the analogues was also tested. At concentrations of 100 nM none of compounds 14, 1, 15, 5x, 6-11, 13, 16 were toxic when only the analogues were put in the cell. When the hippocampal neurons were pretreated with 100 nM of analogue and then exposed to camptothecin, the neuronal survival for compounds 14, 1, 15, 5x and 5 was 61%, 65%, 23%, 29% and 62% respectively. See Figures 4 and 5. Compounds 14 and 1 had about the same neuroprotective activity as
15 PFT α and exhibited significant neuroprotection against the DNA-damage agent camptothecin. Compound 5x, a 5-methyl substituted derivative of compound 5, and compound 15 could not protect neurons effectively against camptothecin (a topoisomerase I inhibitor). When these analogues 14, 1, 15, 5x, 6-11, 13, 16 were tested their ability to prevent neuron against etoposide, compound 14 and compound 1 were
20 more active than compound PFT α , and compound 15 and compound 5x have activity comparable to PFT α . Compounds 14, 1, 15, 5x, and 5 had activity against the topoisomerase II inhibitor, etoposide.

Compound 14 and compound 1 had neuroprotection against death induced by both camptothecin and etoposide. When these two compounds were selected for further
25 study, they were also shown to suppress caspase activation and protect cultured hippocampal neurons against death induced by glutamate. Neuron survival was increased 133% and 39% respectively when neurons were pretreated with compound 14 or 1.

Thus, compounds 14, 1, 15, 5x, 8 and 9 proved to be highly potency in protecting
30 primary hippocampal cells from apoptosis induced cell death. The N-substituent group of compound 5 proved to be essential for biological action. Modification of N-substituent to omit the carbonyl group, compound 14, resulted in a compound that retained biological activity in the protection of hippocampal cells from both etoposide and camptothecin. 4-Methyl substitution in the phenyl position of the N-substituent
35 proved to be non essential for biological action, with compound 13 proving to be highly active in PC12 cells only. Indeed, substitution of a simple alkyl, compound 1, for the

5 aryl group in the N-substituent was well tolerated to provide high activity in hippocampal cells. Substitution of electron donating (e.g., OCH₃ as in compound 6) and withdrawing (e.g., Cl as in compound 9) groups, particularly in the 4'-position of the phenyl group of the N-substituent, were well tolerated. Methyl substitution in position 5, as in compound 5x, was well tolerated and provided an agent active in all assays.

10

Example 6

Neuroprotective Activity of PFT α (2) and its Precursor in an *In Vivo* Stroke Model

15 Surgery was performed in anesthetized (xylazine 5 mg/kg, chloral hydrate 350 mg/kg) mice (3 month old C57Bl/6) in accordance with a protocol approved by the NIH Animal Care and Use Committee. The focal ischemia/reperfusion model has been described previously. See Bruce et al. (1996) *Nat. Med.* 2:788-94. This method involves occluding the middle cerebral artery for 1 h with a nylon thread, and then
20 removing it to allow reperfusion. During the procedure, mice were maintained at 37°C and blood gases, flow and pressure were monitored. At 24 h after ischemia, mice were anesthetized, decapitated, and their brains were cut into 2 mm coronal sections, which then were stained with triphenyltetrazolium for 30 min at 37°C. Images of the stained brains were captured by digital camera for quantitative analysis of infarct area and
25 volume.

PFT α (5) (2 mg/kg, i.p. in 0.2% DMSO) administered 30 min prior to the initiation of cerebral ischemia exhibited neuroprotective activity. Figure 6A shows the infarct area at four different brain levels, with a reduction in the infarct area following PFT α . The infarct volume for vehicle was 40 mm³. When mice were treated with
30 PFT α , infarct volume was 20 mm³, a highly significant 50% decrease. See Figure 6B. Thus, after systemic administration, PFT α entered and acted in the brain.

5

Example 7Neuroprotective Activity of PFT α and its Analogues in an *In Vivo* Model of Parkinson's Disease

The ability of PFT α (5) and compounds 5x (Z-1-117) and 13 (Z-1-143) to
10 protect dopaminergic neurons against MPTP toxicity was tested. MPTP (20 mg/kg)
was administered intraperitoneally to 2 month-old male C57BL/6 mice by four
injections at two hour intervals to induce dopaminergic cell death. Animals were
pretreated with either PFT α (2mg/kg), compound 5x (2mg/kg), compound 13
(2mg/kg), or vehicle (0.2% DMSO) by intraperitoneal injection 30 min. prior to the
15 first MPTP injection and 30 min. after the last MPTP injection.

At seven days, behavioral tests were performed using the rotarod test
equipment. The number of falls and running times were measured. See Figure 7.
Treatment with PFT- α and compound 5x resulted in a statistically significant decrease
in the number of falls caused by MPTP treatment. See Figure 7A. Also, PFT- α and
20 compound 5x resulted in a statistically significant increase in the running time, which is
markedly reduced by MPTP treatment as compared to control. See Figure 7B.

Levels of tyrosine hydroxylase (TH), the enzyme that catalyzes the rate-limiting
step in dopamine synthesis, also were assessed in the striatum using Western blot
analysis. See Figure 8. Densitometric analysis shows that PFT- α and compound 5x
25 reverse the MPTP reduction in TH levels.

In a separate experiment, levels of striatal dopamine, dopamine metabolites,
serotonin, and serotonin metabolites were measured using HPLC analysis.
Specifically, mice were euthanatized, and striatal tissue samples were removed and
levels of dopamine, DOPAC, HVA, 5-HT, 5-HIAA were quantified. The data are
30 shown in Table 2. PFT- α and compound (5x) resulted in a statistically significant
increase in levels of dopamine, DOPAC, and HVA as compared to the MPTP-induced
reduction over control. There was no significant difference in levels of serotonin or
serotonin metabolites.

Table 2

	Dopamine	DOPAC	HVA	5-HT	5-HIAA
Control	6178 ± 799	337 ± 42	618 ± 53	486 ± 33	298 ± 29
MPTP	405 ± 56	undetectable	124 ± 53	441 ± 22	239 ± 24
Vehicle+ MPTP	404 ± 93	undetectable	125 ± 37	406 ± 17	243 ± 18
PFT-α (5)+ MPTP	3314 ± 670**	170 ± 25 **	328 ± 35**	419 ± 20	245 ± 33
Z-1-117 (5x)+ MPTP	1833 ± 256*	102 ± 11 **	302 ± 20**	424 ± 12	246 ± 15
Z-1-143 (13)+ MPTP	440 ± 122	10 ± 10	91 ± 58	429 ± 42	223 ± 45

Values (pg/mg tissue weight) are the mean ± SE of determinations made in 4-6 mice/group.

* $P < 0.05$, ** $P < 0.01$ compared to the value of MPTP group. ANOVA with Scheffe post-hoc tests.

In yet another experiment, the number of TH-immunostained cells in the substantia nigra was assessed in four animals seven days after treatment. Mice were euthanized and perfused by 4% paraformaldehyde transcardially. Brain sections were

- 5 cut and immunostained with an antibody against TH. PFT- α and compound 5x attenuated MPTP-induced loss of substantia nigra dopaminergic neurons. See Figure 9.

Example 8

Neuroprotective Activity of PFT Analogues to an Excitotoxic Insult


10

Primary hippocampal neurons were prepared and cultured as described above. Cultures were pretreated with compounds 14 and 1 (0.04 μ M) 1h before exposure to glutamate (20 μ M) in Locke's medium. The percentage of neuronal survival 24h after the onset of insult is given as mean values \pm SD. Glutamate resulted in a marked decrease in neuronal survival. The glutamate-induced decrease was at least partially reversed by compound 14, thus demonstrating a neuroprotective effect. See Figure 10.

Example 9

Scheme B

20



25 Z-1-169: 2,6-Bis[(4,5,6,7-tetrahydro-2-imino-3(2H)-benzothiazolyl)methyl]pyridine dihydrobromide (30)

This was prepared by a similar procedure from 2-amino-4,5,6,7-tetrahydrobenzothiazole (109 mg, 0.7 mmol) and 2,6-Bis(bromomethyl)pyridine (95 mg, 0.35 mmol); yield (61%); ^1H NMR (DMSO- d_6) δ 9.29 (s, 2H), 7.96 (t, J = 7.7 Hz,

- 5 1H), 7.47 (d, J = 7.7Hz, 2H), 5.33 (s, 4H), 2.56 (m, 4H), 2.12 (m, 4H), 1.68 (m, 8H);
Anal. (C₂₁H₂₇Br₂N₅S₂) C, H, N.

In formula (XV) above, R₁ is H, C₁₋₈ alkyl or aryl, substituted or unsubstituted;
R₂ is any aromatic ring, substituted or unsubstituted and X is S or O.

10


Throughout this application, various publications are referenced. The disclosures of these publications in their entireties are hereby incorporated by reference into this application in order to more fully describe the state of the art to which this invention pertains.

15

- It will be apparent to those skilled in the art that various modifications and variations can be made in the present invention without departing from the scope or spirit of the invention. Other embodiments of the invention will be apparent to those skilled in the art from consideration of the specification and practice of the invention disclosed herein. It is intended that the specification and examples be considered as exemplary only, with a true scope and spirit of the invention being indicated by the following claims.
- 20

What is claimed is:

1. A tetrahydrobenzothiazole analogue having the formula (I):




or a pharmaceutically acceptable salt or ester thereof, wherein:

X is O or S;

Y is NH, O, NR₂ or S;

Z is N or CH;


 = is a mono-substituted, di-substituted, tri-substituted, or quad-substituted 6-member ring that is saturated or unsaturated, wherein the R₁ substituents are independently chosen from H, CN, NO₂, S, N, O, OH, COOH, halogen, and R₂, but not methyl; and

R₂ is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of one carbon atom or from 3 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the

group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;

- (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₂-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy, wherein the halogen is not in the para position;
- (d) bisaryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.
2. A pharmaceutical composition comprised of a compound of claim 1 in combination with a pharmaceutically acceptable carrier.
3. A tetrahydrobenzothiazole analogue having the formula (II):




(II)

or a pharmaceutically acceptable salt or ester thereof, wherein:

X is O or S;

Y is NH, O, NR₂ or S;

Z is N or CH;




= is a mono-substituted, di-substituted, tri-substituted, or quad-substituted 6-member ring that is saturated or unsaturated, wherein the R₁ substituents are independently chosen from H, CN, NO₂, S, N, O, OH, COOH, halogen, and R₂, but not methyl ; and

R₂ is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of one to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;
- (c) aryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
- (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

4. A pharmaceutical composition comprised of a compound of claim 3 in combination with a pharmaceutically acceptable carrier.

5. A tetrahydrobenzothiazole analogue having the formula (III):




or a pharmaceutically acceptable salt or ester thereof, wherein:

R is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of one carbon atom or from 3 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;
- (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₂-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy, wherein the halogen is not in the para position;
- (d) bisaryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl;
- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O,

OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

6. An analogue of claim 5, wherein R is ethyl, methoxy, methoxyphenyl, fluorophenyl, chlorophenyl, nitrophenyl, or tetradecyloxyphenyl.
7. A pharmaceutical composition comprised of a compound of claim 5 in combination with a pharmaceutically acceptable carrier.
8. A tetrahydrobenzothiazole analogue having the formula (IV):




or a pharmaceutically acceptable salt or ester thereof, wherein:

R is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;

- (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
- (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.
9. An analogue of claim 8, wherein R is benzyl, cyclopropylmethyl, fluorobenzyl, difluorobenzyl, methylbenzyl, t-butylbenzyl, nitrobenzyl, cyanobenzyl, trifluoromethylbenzyl, phenylbenzyl, menaphthyl, phenylethyl, or butyl.
10. A pharmaceutical composition comprised of a compound in accordance with claim 8 in combination with a pharmaceutically acceptable carrier.
11. A tetrahydrobenzothiazole analogue having the formula (V):




or a pharmaceutically acceptable salt or ester thereof, wherein:

R is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
 - (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;
 - (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
 - (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
 - (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.
12. An analogue of claim 11, wherein R is methyl, benzyl, cyclopropylmethyl, fluorobenzyl, difluorobenzyl, chlorobenzyl, dichlorobenzyl, methylbenzyl, t-butylbenzyl, nitrobenzyl, cyanobenzyl, trifluoromethylbenzyl, phenylbenzyl, menaphthyl, phenylethyl, or butyl.
13. A pharmaceutical composition comprised of a compound in accordance with claim 11 in combination with a pharmaceutically acceptable carrier.

14. A tetrahydrobenzothiazole analogue having the formula (VI):




or a pharmaceutically acceptable salt or ester thereof, wherein:

R is selected from the group consisting of:

- (a) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
 - (b) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
 - (c) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.
15. An analogue of claim 14, wherein R is methyl, benzyl, cyclopropylmethyl, fluorobenzyl, difluorobenzyl, chlorobenzyl, dichlorobenzyl, methylbenzyl, t-butylbenzyl, nitrobenzyl, cyanobenzyl, trifluoromethylbenzyl, phenylbenzyl, menaphthyl, phenylethyl, or butyl.
16. A pharmaceutical composition comprised of a compound in accordance with claim 14 in combination with a pharmaceutically acceptable carrier.

17. A tetrahydrobenzothiazole analogue having the formula (VII):




or a pharmaceutically acceptable salt or ester thereof, wherein:

R is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;
- (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
- (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

18. An analogue of claim 17, wherein R is methyl, benzyl, cyclopropylmethyl, fluorobenzyl, difluorobenzyl, chlorobenzyl, dichlorobenzyl, methylbenzyl, t-butylbenzyl, nitrobenzyl, cyanobenzyl, trifluoromethylbenzyl, phenylbenzyl, menaphthyl, phenylethyl, or butyl.
19. A pharmaceutical composition comprised of a compound in accordance with claim 17 in combination with a pharmaceutically acceptable carrier.
20. A tetrahydrobenzothiazole analogue having the formula (VIII):




(VIII)

or a pharmaceutically acceptable salt or ester thereof, wherein:

R is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;


- (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
- (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.
21. An analogue of claim 20, wherein R is methyl, benzyl, cyclopropylmethyl, fluorobenzyl, difluorobenzyl, chlorobenzyl, dichlorobenzyl, methylbenzyl, t-butylbenzyl, nitrobenzyl, cyanobenzyl, trifluoromethylbenzyl, phenylbenzyl, menaphthyl, phenylethyl, or butyl.
22. A pharmaceutical composition comprised of a compound in accordance with claim 20 in combination with a pharmaceutically acceptable carrier.
23. A tetrahydrobenzooxazole analogue having the formula (IX):



or a pharmaceutically acceptable salt or ester thereof, wherein:

R is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
 - (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;
 - (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
 - (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
 - (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.
24. An analogue of claim 23, wherein R is methyl, benzyl, cyclopropylmethyl, fluorobenzyl, difluorobenzyl, chlorobenzyl, dichlorobenzyl, methylbenzyl, t-butylbenzyl, nitrobenzyl, cyanobenzyl, trifluoromethylbenzyl, phenylbenzyl, menaphthyl, phenylethyl, or butyl.
25. A pharmaceutical composition comprised of a compound in accordance with claim 23 in combination with a pharmaceutically acceptable carrier.
26. A tetrahydrobenzooxazole analogue having the formula (X):




or a pharmaceutically acceptable salt or ester thereof, wherein:

R is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;
- (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
- (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

27. An analogue of claim 26, wherein R is methyl, benzyl, cyclopropylmethyl, fluorobenzyl, difluorobenzyl, chlorobenzyl, dichlorobenzyl, methylbenzyl, t-butylbenzyl, nitrobenzyl, cyanobenzyl, trifluoromethylbenzyl, phenylbenzyl, menaphthyl, phenylethyl, or butyl.
28. A pharmaceutical composition comprised of a compound in accordance with claim 26 in combination with a pharmaceutically acceptable carrier.
29. A method of reducing or delaying apoptosis in a population of cells, comprising contacting the population of cells with a tetrahydrobenzothiazole analogue, thereby reducing or delaying apoptosis in the population of cells.
30. The method of claim 29, wherein the tetrahydrobenzothiazole analogue comprises pifithrin- α .
31. The method of claim 29, wherein the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (I):




or a pharmaceutically acceptable salt or ester thereof, wherein:

X is O or S;

Y is NH, O, NR₂ or S;

Z is N or CH;




= is a mono-substituted, di-substituted, tri-substituted, or quad-substituted 6-member ring that is saturated or unsaturated, wherein the R₁ substituents are independently chosen from H, CN, NO₂, S, N, O, OH, COOH, halogen, and R₂, but not methyl; and

R₂ is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of one carbon atom or from 3 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;
- (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₂-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy, wherein the halogen is not in the para position;
- (d) bisaryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

32. The method of claim 29, wherein the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (II):




(II)

or a pharmaceutically acceptable salt or ester thereof, wherein:

X is O or S;

Y is NH, O, NR₂ or S;

Z is N or CH;




is a mono-substituted, di-substituted, tri-substituted, or quad-substituted 6-member ring that is saturated or unsaturated, wherein the R₁ substituents are independently chosen from H, CN, NO₂, S, N, O, OH, COOH, halogen, and R₂, but not methyl ; and

R₂ is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of one to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;
- (c) aryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;

- (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

33. The method of claim 29, wherein the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (III):




or a pharmaceutically acceptable salt or ester thereof, wherein:

R is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of one carbon atom or from 3 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;

- (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₂-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy, wherein the halogen is not in the para position;
- (d) bisaryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl;
- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

34. The method of claim 29, wherein the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (IV):



or a pharmaceutically acceptable salt or ester thereof, wherein:


R is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting

of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;

- (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
- (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

35. The method of claim 29, wherein the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (V):




or a pharmaceutically acceptable salt or ester thereof, wherein:

R is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;

- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;
- (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
- (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.
36. The method of claim 29, wherein the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (VI):




(VI)

or a pharmaceutically acceptable salt or ester thereof, wherein:

R is selected from the group consisting of:

- (a) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;

- (b) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
- (c) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.
37. The method of claim 29, wherein the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (VII):




or a pharmaceutically acceptable salt or ester thereof, wherein:

R is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;

- (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
- (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

38. The method of claim 29, wherein the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (VIII):



(VIII)

or a pharmaceutically acceptable salt or ester thereof, wherein:


R is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting

of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;

- (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
- (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

39. The method of claim 29, wherein the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (IX):



(IX)

or a pharmaceutically acceptable salt or ester thereof, wherein:


R is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member

selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;

- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;
- (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
- (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

40. The method of claim 29, wherein the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (X):




or a pharmaceutically acceptable salt or ester thereof, wherein:

R is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
 - (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;
 - (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
 - (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
 - (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.
41. The method of claim 29, wherein the contacting step is *in vivo*.
42. The method of claim 29, wherein the contacting step is *in vitro*.
43. The method of claim 29, wherein the cells are neural cells.
44. The method of claim 43, wherein the neural cells are neuronal cells.
45. The method of claim 29, wherein the cells are cardiac cells.

46. The method of claim 45, wherein the cells are cardiomyocytes.
47. The method of claim 29, wherein the cells are muscle cells.
48. The method of claim 47, wherein the muscle cells are skeletal muscle cells.
49. The method of claim 29, wherein the cells are pancreatic islet cells.
50. The method of claim 29, wherein the apoptosis is induced by a toxin.
51. The method of claim 29, wherein the apoptosis is induced by an environmental factor.
52. The method of claim 29, wherein the apoptosis is induced by a degenerative condition.
53. The method of claim 29, wherein the apoptosis is induced by a severe seizure disorder.
54. The method of claim 29, wherein the apoptosis is induced by a genetic disease.
55. The method of claim 50, wherein the toxin is selected from the group consisting of a neurotoxic form of amyloid β -peptide, camptothecin, glutamate, etoposide, anti-cancer drugs, vinca alkaloids, 3-nitrotyrosine, MPTP, domoic acid, and kainic acid.
56. The method of claim 29, wherein the apoptosis is induced by ischemia.
57. The method of claim 52, wherein the ischemia is induced by a stroke.
58. The method of claim 42, wherein the ischemia is induced by a myocardial infarction.

59. The method of claim 29, wherein the apoptosis is induced by trauma.
60. The method of claim 29, wherein the apoptosis is induced by a genetic defect.
61. A method of treating a subject with a degenerative condition or of reducing one or more symptoms of a degenerative condition in a subject, comprising administering to the subject a therapeutically effective amount of a tetrahydrobenzothiazole analogue.
62. The method of claim 57, wherein the tetrahydrobenzothiazole analogue comprises pifithrin- α .
63. The method of claim 57, wherein the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (I):




or a pharmaceutically acceptable salt or ester thereof, wherein:

X is O or S;

Y is NH, O, NR₂ or S;

Z is N or CH;




= is a mono-substituted, di-substituted, tri-substituted, or quad-substituted 6-member ring that is saturated or unsaturated, wherein the R₁ substituents are independently chosen from H, CN, NO₂, S, N, O, OH, COOH, halogen, and R₂, but not methyl ; and

R₂ is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of one carbon atom or from 3 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;
- (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₂-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy, wherein the halogen is not in the para position;
- (d) bisaryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

64. The method of claim 57, wherein the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (II):




or a pharmaceutically acceptable salt or ester thereof, wherein:

X is O or S;

Y is NH, O, NR₂ or S;

Z is N or CH;




= is a mono-substituted, di-substituted, tri-substituted, or quad-substituted 6-member ring that is saturated or unsaturated, wherein the R₁ substituents are independently chosen from H, CN, NO₂, S, N, O, OH, COOH, halogen, and R₂, but not methyl ; and

R₂ is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of one to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;

- (c) aryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
- (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

65. The method of claim 57, wherein the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (III):




or a pharmaceutically acceptable salt or ester thereof, wherein:

R is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;

- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of one carbon atom or from 3 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;
- (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₂-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy, wherein the halogen is not in the para position;
- (d) bisaryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl;
- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

66. The method of claim 57, wherein the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (IV):



or a pharmaceutically acceptable salt or ester thereof, wherein:


R is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member

selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;

- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;
- (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
- (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.


67. The method of claim 57, wherein the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (V):



or a pharmaceutically acceptable salt or ester thereof, wherein:
R is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;
- (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
- (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

68. The method of claim 57, wherein the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (VI):




or a pharmaceutically acceptable salt or ester thereof, wherein:

R is selected from the group consisting of:

- (a) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
- (b) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
- (c) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

69. The method of claim 57, wherein the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (VII):




or a pharmaceutically acceptable salt or ester thereof, wherein:

R is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;
- (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
- (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and

- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

70. The method of claim 57, wherein the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (VIII):



or a pharmaceutically acceptable salt or ester thereof, wherein:


R is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;
- (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
- (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH,

halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and

- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

71. The method of claim 57, wherein the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (IX):




or a pharmaceutically acceptable salt or ester thereof, wherein:

R is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;
- (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;

- (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

72. The method of claim 57, wherein the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (X):



or a pharmaceutically acceptable salt or ester thereof, wherein:


R is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;

- (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
 - (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
 - (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.
73. The method of claim 57, wherein the degenerative condition is a neurodegenerative condition.
74. The method of claim 69, wherein the neurodegenerative condition is selected from the group consisting of Alzheimer's disease, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, stroke, multiple sclerosis, brain injury, spinal cord injury, and peripheral neuropathy.
75. The method of claim 57, wherein the degenerative condition is a degenerative cardiomyopathy.
76. The method of claim 57, wherein the degenerative condition is a degenerative myopathy.
77. The method of claim 57, wherein the degenerative condition is diabetes.
78. The method of claim 57, wherein the subject is a human.
79. A method of treating a subject after an ischemic event to reduce ischemia-induced apoptosis, comprising administering to the subject a therapeutically

effective amount of a tetrahydrobenzothiazole analogue, thereby reducing apoptosis.

80. The method of claim 75, wherein the tetrahydrobenzothiazole analogue comprises pifithrin- α .
81. The method of claim 75, wherein the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (I):




(I)

or a pharmaceutically acceptable salt or ester thereof, wherein:

X is O or S;

Y is NH, O, NR₂ or S;

Z is N or CH;



= is a mono-substituted, di-substituted, tri-substituted, or quad-substituted 6-member ring that is saturated or unsaturated, wherein the R₁ substituents are independently chosen from H, CN, NO₂, S, N, O, OH, COOH, halogen, and R₂, but not methyl; and


R₂ is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member

selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;

- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of one carbon atom or from 3 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;
- (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₂-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy, wherein the halogen is not in the para position;
- (d) bisaryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

82. The method of claim 75, wherein the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (II):




or a pharmaceutically acceptable salt or ester thereof, wherein:

X is O or S;

Y is NH, O, NR₂ or S;

Z is N or CH;




= is a mono-substituted, di-substituted, tri-substituted, or quad-substituted 6-member ring that is saturated or unsaturated, wherein the R₁ substituents are independently chosen from H, CN, NO₂, S, N, O, OH, COOH, halogen, and R₂, but not methyl ; and

R₂ is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of one to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;
- (c) aryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
- (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

83. The method of claim 75, wherein the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (III):




or a pharmaceutically acceptable salt or ester thereof, wherein:

R is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of one carbon atom or from 3 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;
- (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₂-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy, wherein the halogen is not in the para position;
- (d) bisaryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl;
- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O,

OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

84. The method of claim 75, wherein the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (IV):




or a pharmaceutically acceptable salt or ester thereof, wherein:

R is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;
- (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
- (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and

- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.
85. The method of claim 75, wherein the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (V):



or a pharmaceutically acceptable salt or ester thereof, wherein:


R is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;
- (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
- (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH,

halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and

- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

86. The method of claim 75, wherein the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (VI):




or a pharmaceutically acceptable salt or ester thereof, wherein:

R is selected from the group consisting of:

- (a) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
- (b) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
- (c) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

87. The method of claim 75, wherein the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (VII):



(VII)


or a pharmaceutically acceptable salt or ester thereof, wherein:

R is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;
- (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
- (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and

- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

88. The method of claim 75, wherein the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (VIII):



or a pharmaceutically acceptable salt or ester thereof, wherein:


R is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;
- (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
- (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH,

halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and

- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

89. The method of claim 75, wherein the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (IX):



(IX)


or a pharmaceutically acceptable salt or ester thereof, wherein:

R is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;

- (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
- (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

90. The method of claim 75, wherein the tetrahydrobenzothiazole analogue comprises one or more analogues of formula (X):



or a pharmaceutically acceptable salt or ester thereof, wherein:

R is selected from the group consisting of:

- (a) alkyl, or alkenyl, or alkynyl, each of which are straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, OH, COOH, and halogen;
- (b) alkoxy which is straight chain, branched chain or cyclic, having a carbon chain length of from 1 to 20 carbon atoms, unsubstituted or substituted with at least one member selected from the group consisting

of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, and C₁-C₂₀ alkynyl;


- (c) aryl which is substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, and C₁-C₂₀ alkoxy;
- (d) bisaryl which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl; and
- (e) condensed aromatic which is unsubstituted or substituted with at least one member selected from the group consisting of CN, NO₂, S, N, O, OH, COOH, halogen, C₁-C₂₀ alkyl, C₁-C₂₀ alkenyl, C₁-C₂₀ alkynyl, C₁-C₂₀ alkoxy, and aryl.

- 91. The method of claim 75, wherein the ischemic event is a stroke.
- 92. The method of claim 75, wherein the ischemic event is a myocardial infarction.
- 93. The method of claim 75, wherein the subject is a human.
- 94. A tetrahydrobenzothiazole analogue wherein the analogue is
 Ethyl 2-(4,5,6,7-tetrahydro-2-imino-3(2*H*)-benzothiazolyl)-acetate
 hydrobromide,
 3-(phenylmethyl)-4,5,6,7-tetrahydro-2(3*H*)-benzothiazolimine hydrobromide,
 1-(4,5,6,7-tetrahydro-2-imino-3(2*H*)-benzothiazolyl)-2-butanone hydrobromide,
 3-(cyclopropylmethyl)-4,5,6,7-tetrahydro-2(3*H*)-benzothiazolimine
 hydrobromide,
 1-(4-methylphenyl)-2-(4,5,6,7-tetrahydro-2-imino-6-methyl-3(2*H*)-
 benzothiazolyl)-ethanone hydrobromide,
 1-(4-methylphenyl)-2-(4,5,6,7-tetrahydro-2-imino-6-methyl-3(2*H*)-
 benzothiazolyl)-ethanone hydrobromide,

1-(2-methoxyphenyl)-2-(4,5,6,7-tetrahydro-2-imino-3(2H)-benzothiazolyl)-ethanone hydrobromide,
1-(3-methoxyphenyl)-2-(4,5,6,7-tetrahydro-2-imino-3(2H)-benzothiazolyl)-ethanone hydrobromide,
1-phenyl-2-(4,5,6,7-tetrahydro-2-imino-3(2H)-benzothiazolyl)-ethanone hydrobromide, or
1-(4-chlorophenyl)-2-(4,5,6,7-tetrahydro-2-imino-3(2H)-benzothiazolyl)-ethanone hydrobromide.

95. A pharmaceutical composition comprised of a compound in accordance with claim 90 in combination with a pharmaceutically acceptable carrier.


96. A method of treating a subject with a degenerative condition or of reducing one or more symptoms of a degenerative condition in a subject, comprising administering to the subject a therapeutically effective amount of a tetrahydrobenzothiazole analogue of the formula XIV



or a pharmaceutically acceptable salt or ester thereof, wherein

R₁ is H, or CH₃, R₂ is C₁₋₈ alkyl or aryl substituted or unsubstituted and X is S or O.

97. A tetrahydrobenzothiazole analogue having the formula (XV)



or a pharmaceutically acceptable salt or ester thereof, wherein

R_1 is H, C1-8 alkyl or aryl substituted or unsubstituted, R_2 is any aromatic ring substituted or unsubstituted and X is S or O.

98. A pharmaceutical composition comprised of a compound of claim 97 in combination with a pharmaceutical acceptable carrier.
99. A method of treating a subject with a degenerative condition or of reducing one or more symptoms of a degenerative condition in a subject, comprising administering to the subject a therapeutically effective amount of the tetrahydrobenzothiazole analogue of claim 97.
100. A method of treating a subject exposed to irradiation, comprising administering to the subject a therapeutically effective amount of a tetrahydrobenzothiazole analogue






	Ref. No.	Structure	Mol. Wt.	¹ H-NMR ¹³ C NMR	m.p.	Solubility
	Z-1-073		367.3049			
1	Z-1-110		305.2356	Z-1-110		Water
15	Z-1-113		289.2362	Z-1-113		Water
14	Z-1-119		325.2683	Z-1-119		Water
5X	Z-1-117		381.3315	Z-1-117		Hot water

Fig. 1




Fig. 2a

40 μ M Camptothecin Challenge Test of PFTalpha Derivatives in PC12 Cells




Fig. 2b




Fig. 3




Fig. 4




Fig. 5

7/11




Fig. 6a




Fig. 6b

8/11




Fig. 7a




Fig. 7b




Fig. 8




Fig. 9




Fig. 10

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
17 January 2002 (17.01.2002)

PCT

(10) International Publication Number
WO 02/004409 A3

(51) International Patent Classification⁷: C07D 277/82,
277/68, 277/72, 263/58, 513/04, 498/04, A61K 31/428,
31/423, A61P 25/00, 25/16, 25/28 // (C07D 513/04,
277:00, 235:00) (C07D 498/04, 263:00, 235:00)

[CN/US]; 116 Ridgefield, Lutherville, MD 21093 (US).
HOLLOWAY, Harold, Wayne [US/US]; 121 Mariners
Point Dr., Middle River, MD 21220 (US).

(21) International Application Number: PCT/US01/21504

(74) Agents: MILLER, Mary, L. et al.: Needle & Rosenberg,
P.C., Suite 1200, The Candler Building, 127 Peachtree
Street, N.E., Atlanta, GA 30303-1811 (US).

(22) International Filing Date: 6 July 2001 (06.07.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/216,388 6 July 2000 (06.07.2000) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK,
SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA,
ZW.

(63) Related by continuation (CON) or continuation-in-part
(CIP) to earlier application:
US 60/216,388 (CIP)
Filed on 6 July 2000 (06.07.2000)

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant and

(72) Inventor (*for all designated States except US*): THE
GOVERNMENT OF THE UNITED STATES OF
AMERICA, as represented by THE SECRETARY,
DEPARTMENT OF HEALTH AND HUMAN SER-
VICES [US/US]; c/o National Institutes of Health, Office
of Technology Transfer, Suite 325, 6011 Executive Boule-
vard, Rockville, MD 20852-3804 (US).

Published:

— with international search report
— before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): GREIG, Nigel, H.
[US/US]; 11 Anne Brent Garth, Phoenix, MD 21131 (US).
MATTSON, Mark [US/US]; 858 E. Wheel Road, Bel Air,
MD 21015 (US). ZHU, Xiaoxiang [US/US]; 6114 East
Pratt Street, Baltimore, MD 21224 (US). YU, Qian-Sheng

(88) Date of publication of the international search report:
18 July 2002

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: TETRAHYDROBENZOTHAZOLE ANALOGUES AS NEUROPROTECTIVE AGENTS

(57) Abstract: This invention relates generally to tetrahydrobenzothiazole analogues and tetrahydrobenzooxazole analogues, to pharmaceutical compositions which include these compounds and a pharmaceutically acceptable carrier, and to methods of treatment using these compounds. The invention also encompasses pharmaceutically acceptable esters, amides, and salts of such compounds. The invention further provides a method of reducing or delaying apoptosis in a population of cells, comprising contacting the population of cells with a tetrahydrobenzothiazole analogue or a tetrahydrobenzooxazole analogue, thereby reducing or delaying apoptosis in the population of cells.

WO 02/004409 A3

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/21504

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D277/82 C07D277/68 C07D277/72 C07D263/58 C07D513/04
C07D498/04 A61K31/428 A61K31/423 A61P25/00 A61P25/16
A61P25/28 //(C07D513/04,277:00,235:00),(C07D498/04,263:00,

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, PAJ, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	KOMAROV P G ET AL: "A CHEMICAL INHIBITOR OF P53 THAT PROTECTS MICE FROM THE SIDE EFFECTS OF CANCER THERAPY" SCIENCE, AMERICAN ASSOCIATION FOR THE ADVANCEMENT OF SCIENCE,, US, vol. 285, no. 5434, 10 September 1999 (1999-09-10), pages 1733-1737, XP000953397 ISSN: 0036-8075 cited in the application page 1733, abstract; page 1734, column 1, line 3 to page 1735, column 3, line 4; page 1735, column 3, last line to page 1737, column 1, line 11; page 1737, column 1, bridge paragraph to column 2 --- -/--	1-95,100



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

3 May 2002

Date of mailing of the international search report

22/05/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Hass, C

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/21504

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 235:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 186 087 A (THOMAE GMBH DR K) 2 July 1986 (1986-07-02) claims 1,8-11	61
X	DE 39 23 689 A (BOEHRINGER INGELHEIM KG) 24 January 1991 (1991-01-24) page 3, line 48 - line 56; claims 1,6	61
X	US 3 462 448 A (KELYMAN JACQUELINE S) 19 August 1969 (1969-08-19) column 2, line 71 -column 3, line 25; claims 1,3	61
X	DE 19 26 558 A (DR. KARL THOMAE GMBH) 3 December 1970 (1970-12-03) claims 1,7,8	61

-/--



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

3 May 2002

Date of mailing of the international search report

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040. Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Hass, C

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/21504

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DE 42 41 013 A (BOEHRINGER INGELHEIM KG) 9 June 1994 (1994-06-09) claims	61
X	PATENT ABSTRACTS OF JAPAN vol. 1998, no. 11, 30 September 1998 (1998-09-30) & JP 10 152488 A (TOKYO TANABE CO LTD), 9 June 1998 (1998-06-09) abstract	61
X	PATENT ABSTRACTS OF JAPAN vol. 1996, no. 09, 30 September 1996 (1996-09-30) & JP 08 113570 A (SANKYO CO LTD; SANKYO KASEI KOGYO KK), 7 May 1996 (1996-05-07) abstract	61
X	US 4 769 370 A (WOLTERSDORF JR OTTO W ET AL) 6 September 1988 (1988-09-06) abstract	61
A	EP 0 705 832 A (LILLY INDUSTRIES LTD) 10 April 1996 (1996-04-10) page 8, line 15 - line 24; claims 1,7,8	1-100
A	EP 0 281 261 A (LUNDBECK & CO AS H) 7 September 1988 (1988-09-07) page 2, line 1 - line 5; claims 1,4	1-100
A	US 4 377 695 A (LAUTENSCHLAEGER HANS-HEINER ET AL) 22 March 1983 (1983-03-22) column 1, line 15 - line 63	1-100
A	US 4 562 190 A (UEDA IKUO ET AL) 31 December 1985 (1985-12-31) claims 1,6,7	1-100
A	US 4 797 399 A (UEDA IKUO ET AL) 10 January 1989 (1989-01-10) claims 1,5,6	1-100
A	US 4 859 700 A (LAVIELLE GILBERT ET AL) 22 August 1989 (1989-08-22) claims 1,7,8	1-100
A	US 5 132 305 A (BONNET JACQUELINE ET AL) 21 July 1992 (1992-07-21) claims 1,8-10	1-100
	-/--	

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/21504

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>SINGH ET AL: "Heterocyclic systems containing bridgehead nitrogen atom: Part XXV-syntheses of imidazo(2,1-b)benzothiazoles and quinoxalino(2,3:4',5')imidazo(2',1'-b)benzothiazoles"</p> <p>INDIAN JOURNAL OF CHEMISTRY, SECTION B: ORGANIC, INCL. MEDICINAL, PUBLICATIONS & INFORMATIONS DIRECTORATE, NEW DELHI, IN, vol. 14B, no. 12, 1976, pages 997-998, XP000953345</p> <p>ISSN: 0019-5103</p> <p>cited in the application table 1</p>	1,5
A	<p>-----</p> <p>CHEMICAL ABSTRACTS, vol. 68, no. 1, 1 January 1968 (1968-01-01)</p> <p>Columbus, Ohio, US;</p> <p>abstract no. 2856m,</p> <p>page 269;</p> <p>XP002198025</p> <p>cited in the application formula (V)</p> <p>& N SALDABOLS ET AL: "Synthesis in the methyl 2-furyl ketone series. XII. 5-Nitro-2-furyl substituted imidazo heterocyclic compounds with a common nitrogen atom"</p> <p>KHIM.-FARM. ZH.,</p> <p>vol. 1, no. 2, 1967, pages 27-31,</p> <p>-----</p>	96

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1, 2, 5-7, 11-13, 17-19, 23-25, 29 (part), 30, 31, 33, 35, 37, 39, 41-61 (part), 62, 63, 65, 67, 69, 71, 73-79 (part), 80, 81, 83, 85, 87, 89, 91-94 (part), 100 (part)

Compounds of formula (I) of claim 1 wherein the alkyl substituent at Z has a keto group in beta-position, and the pharmaceutical use of these compounds.

2. Claims: 3, 4, 8-10, 14-16, 20-22, 26-28, 29 (part), 32, 34, 36, 38, 40, 41-61 (part), 64, 66, 68, 70, 72, 73-79 (part), 82, 84, 86, 88, 90, 91-94 (part), 95, 100 (part)

Compounds of formula (II) of claim 3 wherein the alkyl substituent at Z has no keto group in beta-position, and the pharmaceutical use of these compounds.

3. Claim : 96

Method of medical treatment making use of the tricyclic compound of formula (XIV)

4. Claims: 97-99

Compounds of formula (XV) and their pharmaceutical use.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 29, 41-61, 73, 75-79, 91-93, 100

Present claims 29, 41-61, 73, 75-79, 91-93 and 100 relate to an extremely large number of possibilities. The methods of treatment claimed therein refer to any "tetrahydrobenzothiazole analogues". Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compounds concerned: On page 20 of the description, the expression "tetrahydrobenzothiazole analogue" is defined as to comprise compounds falling actually within the formula (I) of claim 1 and the formula (II) of claim 3. Thus in the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely those parts relating to the compounds comprised by formula (I) of claim 1 and formula (II) of claim 3 and for methods making use of these compounds.

It is noted that all preparative examples and all pharmacological test examples refer to compounds which are in fact derivatives of tetrahydrobenzothiazoles; according to formulae (I) and (II), the claimed subject-matter also comprises the isosteric tetrahydrobenzoxazoles, tetrahydrobenzothiophenes and tetrahydrobenzofurans. All these possibilities have not been verified, however, they may be considered as to represent reasonable generalisations of the very experimental results put forward, and they have been searched as well.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/21504

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0186087	A	02-07-1986	DE 3447075 A1	03-07-1986
			DE 3508947 A1	18-09-1986
			AT 45735 T	15-09-1989
			AU 583874 B2	11-05-1989
			AU 5154485 A	17-07-1986
			BG 62023 B2	30-12-1998
			BR 1100678 A3	13-10-1999
			CA 1263653 A1	05-12-1989
			CS 9104099 A3	16-09-1992
			DD 242230 A5	21-01-1987
			DE 3572485 D1	28-09-1989
			DK 590285 A	23-06-1986
			EP 0186087 A1	02-07-1986
			ES 550235 D0	16-01-1987
			ES 8702787 A1	01-04-1987
			ES 556873 D0	01-08-1987
			ES 8707513 A1	16-10-1987
			ES 556874 D0	01-08-1987
			ES 8707514 A1	16-10-1987
			ES 556875 D0	01-08-1987
			ES 8707515 A1	16-10-1987
			FI 855102 A ,B,	23-06-1986
			GR 853126 A1	22-04-1986
			HK 78692 A	23-10-1992
			HU 39736 A2	29-10-1986
			IE 58863 B	17-11-1993
			IL 77415 A	19-03-1990
			JP 1854941 C	07-07-1994
			JP 61155377 A	15-07-1986
			JP 5072907 B	13-10-1993
			KR 9309791 B1	11-10-1993
			LU 90208 A9	06-04-1998
			MX 9202792 A1	30-06-1992
			NO 855195 A ,B,	23-06-1986
			NZ 214661 A	26-04-1990
			PH 24533 A	03-08-1990
			PT 81735 A ,B	01-01-1986
			SG 82492 G	04-12-1992
			US 4843086 A	27-06-1989
			US 4886812 A	12-12-1989
			US 4731374 A	15-03-1988
			ZA 8509731 A	26-08-1987
DE 3923689	A	24-01-1991	DE 3923689 A1	24-01-1991
			AU 6149690 A	22-02-1991
			CA 2063988 A1	19-01-1991
			DD 298395 A5	20-02-1992
			WO 9101136 A1	07-02-1991
			EP 0483252 A1	06-05-1992
			HU 60922 A2	30-11-1992
US 3462448	A	19-08-1969	NONE	
DE 1926558	A	03-12-1970	DE 1926558 A1	03-12-1970
DE 4241013	A	09-06-1994	DE 4241013 A1	09-06-1994
			AT 183091 T	15-08-1999
			AU 674378 B2	19-12-1996

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/21504

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 4241013	A	AU 5715294 A	04-07-1994
		BG 62475 B1	30-12-1999
		BG 99700 A	30-04-1996
		CA 2150131 A1	23-06-1994
		CN 1091635 A ,B	07-09-1994
		CZ 9501440 A3	13-03-1996
		DE 59309733 D1	16-09-1999
		DK 675717 T3	28-02-2000
		WO 9413287 A1	23-06-1994
		EP 0675717 A1	11-10-1995
		ES 2136725 T3	01-12-1999
		FI 952734 A	05-06-1995
		GR 3031521 T3	31-01-2000
		HK 1019003 A1	08-09-2000
		HU 71798 A2	28-02-1996
		IL 107870 A	05-12-1996
		JP 8504201 T	07-05-1996
		JP 3221880 B2	22-10-2001
		MX 9307644 A1	30-06-1994
		NO 952207 A	02-06-1995
		NZ 258876 A	27-07-1997
		PL 309282 A1	02-10-1995
		RU 2157685 C2	20-10-2000
		SG 49228 A1	18-05-1998
		SK 74095 A3	08-05-1996
		US 6191153 B1	20-02-2001
		ZA 9309065 A	04-08-1994
JP 10152488	A	09-06-1998	NONE
JP 08113570	A	07-05-1996	NONE
US 4769370	A	06-09-1988	NONE
EP 0705832	A	10-04-1996	
		AU 698580 B2	05-11-1998
		AU 3049795 A	28-03-1996
		CA 2157998 A1	13-03-1996
		CN 1129219 A ,B	21-08-1996
		CZ 9502322 A3	13-03-1996
		EP 0705832 A1	10-04-1996
		FI 954243 A	13-03-1996
		HU 72593 A2	28-05-1996
		IL 115236 A	15-06-1998
		JP 8081464 A	26-03-1996
		NO 953575 A	13-03-1996
		NZ 272961 A	26-02-1998
		PL 310373 A1	18-03-1996
		RU 2146256 C1	10-03-2000
		TR 960794 A2	21-10-1996
		US 5563147 A	08-10-1996
		ZA 9507607 A	17-05-1996
EP 0281261	A	07-09-1988	
		AT 75723 T	15-05-1992
		AU 608293 B2	28-03-1991
		AU 1216788 A	01-09-1988
		CA 1338934 A1	25-02-1997
		DE 3870666 D1	11-06-1992
		DK 91488 A	27-08-1988

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/21504

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0281261 A		EP 0281261 A2	07-09-1988
		ES 2037208 T3	16-06-1993
		FI 880906 A ,B,	27-08-1988
		GR 3004984 T3	28-04-1993
		IE 61257 B	19-10-1994
		IL 85342 A	14-01-1993
		JP 2677588 B2	17-11-1997
		JP 63264557 A	01-11-1988
		NO 880833 A ,B,	29-08-1988
		NZ 223440 A	26-02-1991
		PT 86836 A ,B	01-03-1988
		US 4847254 A	11-07-1989
		US 4946863 A	07-08-1990
		ZA 8801217 A	25-01-1989
US 4377695 A	22-03-1983	DE 3042481 A1	16-06-1982
		DK 495881 A	12-05-1982
		EP 0051827 A1	19-05-1982
		GR 74705 A1	05-07-1984
		JP 57108074 A	05-07-1982
		ZA 8107697 A	27-10-1982
US 4562190 A	31-12-1985	AT 37368 T	15-10-1988
		DE 3474145 D1	27-10-1988
		EP 0132817 A1	13-02-1985
		JP 1788070 C	10-09-1993
		JP 4081591 B	24-12-1992
		JP 60034964 A	22-02-1985
US 4797399 A	10-01-1989	AU 6762687 A	23-07-1987
		CN 87100707 A	02-09-1987
		DK 8687 A	18-07-1987
		EP 0232740 A1	19-08-1987
		FI 865369 A	18-07-1987
		JP 62181253 A	08-08-1987
		NO 870053 A	20-07-1987
		OA 8462 A	29-07-1988
		PT 84103 A	01-02-1987
		ZA 8609748 A	26-08-1987
US 4859700 A	22-08-1989	FR 2611713 A1	09-09-1988
		AT 63309 T	15-05-1991
		AU 598345 B2	21-06-1990
		AU 1234488 A	01-09-1988
		CA 1313871 A1	23-02-1993
		DE 3862660 D1	13-06-1991
		DK 104388 A	28-08-1988
		EP 0282390 A1	14-09-1988
		ES 2009553 A6	01-10-1989
		GR 88100111 A ,B	16-12-1988
		IE 60500 B	27-07-1994
		JP 1845506 C	25-05-1994
		JP 5058633 B	27-08-1993
		JP 63227582 A	21-09-1988
		NZ 223663 A	29-01-1990
		OA 8813 A	31-03-1989
		PT 86851 A ,B	01-03-1988
		US 4885299 A	05-12-1989

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/21504

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4859700	A	ZA 8801383 A	28-12-1988
US 5132305	A	21-07-1992	
		FR 2643634 A1	31-08-1990
		AU 626080 B2	23-07-1992
		AU 5019690 A	06-09-1990
		CA 2010989 A1	31-08-1990
		EP 0385848 A1	05-09-1990
		JP 2268172 A	01-11-1990
		JP 6002753 B	12-01-1994
		NZ 232701 A	29-01-1991
		OA 9439 A	15-10-1992
		PT 93278 A	31-08-1990
		US 5182278 A	26-01-1993
		ZA 9001543 A	30-01-1991

THIS PAGE BLANK (USPTO)